

## Short Communication

# Detection of Human Coronavirus NL63 in a 28 Days Old Newborn in Iran

Sultani M<sup>1</sup>, Mokhtari-Azad T<sup>1</sup>, Yavarian J<sup>1\*</sup>

1. Virology Department, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

Respiratory infections are one of the most frequent diseases with high morbidity and mortality in the first years of life. Human coronaviruses 229E, OC43, HKU1 and NL63 are known to cause common cold and upper respiratory tract infections worldwide. Recently it was shown that they can cause severe respiratory disease especially in neonates and infants (1). In this report we collected 322 respiratory tract swabs and washes from children less than two years old who were hospitalized with influenza like illnesses during January to March 2012 in Tehran and surrounded cities. All samples were subjected to RNA extraction with High Pure Viral Nucleic Acid Kit (Roche, Germany). At first Real time RT-PCR was performed for detection of Influenza A and B viruses. Then multiplex SYBR Green Real time RT-PCR was performed for detection of human parainfluenza viruses 1-4, Human coronaviruses 229E, OC43, HKU1 and NL63, rhinoviruses, human metapneumovirus, enteroviruses and respiratory syncytial virus (RSV). In 322 samples, 26 (8%) influenza A/H3N2, 5 (1.5%) influenza B, 22 (6.8%) RSV, 2 (0.6%) parainfluenza virus 3, 2 (0.6%) parainfluenza virus 4 and 1 (0.3%) human coronavirus NL63 (2). Conventional RT-PCR was performed on the coronavirus NL63 positive sample and the product was sequenced. Although nucleotide analysis

confirmed the presence of NL63 genome in the sample, since the length of product was less than 200 nts, it has not been submitted in GenBank.

This was the first detection of coronavirus NL63 in Iran. The patient was a 28 days old newborn girl who was hospitalized with respiratory distress, apnea, cough, stuffed nose, cyanosis and vomiting. She was admitted to NICU and her condition was poor. Her blood culture was negative and all other tests were at the normal range. She passed away because of the respiratory distress.

Here we investigated the viral etiology of respiratory tract infections in 322 children less than two years of age who were hospitalized from January to March 2012 in Tehran. Two principal pathogens were influenza A/H3N2 and RSV the same as many other studies (3). We detected coronavirus NL63 in a newborn with acute respiratory infection which shows that human coronaviruses can lead to the severe respiratory infections and hospitalization. According to the result of this study in addition to influenza virus and RSV, the evaluation of human coronavirus infections is recommended for children hospitalized with influenza like illnesses.

## Conflict of interest

None

## Acknowledgments

The authors thank the entire staff of the National Influenza Center, School of Public Health, Tehran University of Medical

\*Corresponding author: Jila Yavarian, Ph.D. Virology Department, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran, Telefax: (+98) 21 88962343. Email: yavarian@tums.ac.ir

## Human CoV-NL63 in Iran

Sciences. All phases of this study were supported by Tehran University of Medical Sciences grant, No. 91-02-27-17902.

### References

1. Lu S, Wang S, Zhang L, Xu C, Bian C, Wang Z, et al. Epidemiology of Human Respiratory Viruses in Children with Acute Respiratory Tract Infections in Jinan, China. *Clin Dev Immunol*. 2013.
2. Sultani M, Mokhtari Azad T, Eshragian MR, Shadab A, Naseri M, Eilami O, Yavarian J. Multiplex SYBR Green Real time PCR assay for detection of respiratory viruses. *Jundishapur J Microbiol*. In press 2015.
3. Malekshahi SS, Azad TM, Yavarian J, Shahmahmoodi S, Naseri M, Rezaei F. Molecular detection of respiratory viruses in clinical specimens from children with acute respiratory disease in Iran. *Pediatr Infect Dis J*. 2010;29:931-3.