# **Review Article**

# The Worldwide Epidemiology of Hepatitis B, Hepatitis C, and Herpes Simplex Virus Type 2 in Female Sex Workers: A Systematic Review and Meta-Analysis

Farahmand M<sup>1</sup>, Khales P<sup>2</sup>, Monavari SH<sup>2</sup>, Kiani SJ<sup>2</sup>, Tavakoli A<sup>2,3\*</sup>

- 1. Department of Virology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
- 2. Department of Medical Virology, Faculty of Medicine, Iran University of Medical Sciences, Tehran, Iran
- 3. Research Center of Pediatric Infectious Diseases, Institute of Immunology and Infectious Diseases, Iran University of Medical Sciences, Tehran, Iran

#### **Abstract**

**Background and Aims:** Female sex workers are vulnerable and at high risk to acquire sexually transmitted infections, and act as a bridge in the transmission of sexually transmitted infections (STIs) to the general population. This study was conducted to determine the prevalence of the main viral pathogens responsible for STIs, including Hepatitis B virus (HBV), Hepatitis C virus (HCV), and Herpes simplex virus type 2 (HSV-2) in female sex workers in the world.

**Materials and Methods:** A systematic search was carried out for relevant literature in international databases from database inception to September 25, 2019. The pooled prevalence for each STI of interest was estimated by a DerSimonian-Laird random-effects model using the inverse variance method.

**Results:** The lowest pooled prevalence of HBV, HCV, and HSV-2 infections was seen in the Republic of Mauritius (0.17%; 95% CI: 0.01%-2.64%), Panama (0.20%; 95% CI: 0.05%-0.80%), and Iran (14.12%; 95% CI: 6.66%-27.48%), respectively. The highest pooled prevalence of HBV infection was found in Slovakia (22.22%; 95% CI: 8.60%-46.47%), while for HCV and HSV-2 infections was observed in Scotland (64.29%; 95% CI: 54.35%-73.13%) and Indonesia (90.30%; 95% CI: 84.76%-93.97%), respectively.

**Conclusion:** A decreasing trend was observed in the prevalence of STIs of interest among female sex workers during recent years. However, the prevalence has remained high in some regions, and therefore it is important to improve prevention programs and conduct surveillance regularly in all parts of the world to decrease the risk of transmission of infections to the general population.

**Keywords:** Female sex worker; Prostitute; sexually transmitted infection; Hepatitis; Herpes simplex virus; Meta-Analysis

#### Introduction

emale sex workers are defined as women who offer sexual services in return for money, goods, or other markers of economic remuneration. However, female sex workers are a heterogeneous population group who work in varied work environments and community organizations. Street-based sex workers are mainly illegal workers who solicit

\*Corresponding author:

Ahmad Tavakoli.

Email: Tavakoli.a@iums.ac.ir.

clients on the street or in public settings such as markets, parks, cinema halls, and service them in alleys, or the clients' cars. Another group is indoor-based sex workers who are employed to work in brothels, hotels, massage parlors, saunas, and lodges (1-3).

Female sex workers are known to be a highrisk group for the acquisition of sexually transmitted infections (STIs) due to the risk factors associated with their occupation, such as multiple sexual partners, low adherence to condom use, and unsafe sexual practices. This population has long been recognized as a major driver of growing epidemics of STIs in several countries. These epidemics are initially triggered by female sex workers and subsequently generalized to their clients and partners (4, 5).

To date, there are more than 30 different sexually transmissible agents, including viruses, bacteria, and parasites. Among them, three bacterial (Chlamydia trachomatis, Neisseria gonorrhoeae, Treponema pallidum), a parasitic (Trichomonas vaginalis), and six viral (Hepatitis B virus (HBV), Hepatitis C virus (HCV), Human immunodeficiency viruses (HIV), Herpes simplex virus type 2 (HSV-2), and Human papillomavirus (HPV) and Human T-cell leukemia virus type 1 (HTLV-1) infections account for the majority of STIs globally (6, 7).

According to the last reports of the world health organization (WHO), HIV, HBV and HCV are major public health concerns which have significant impacts on human health and human life. It is estimated that approximately 38 million are HIV-infected (8), 257 million are HBV-infected (9), and 71 million have chronic HCV infections worldwide (10). It has been well documented that persistent infections with HCV and HBV are a leading cause of cirrhosis and hepatocellular carcinoma (11, 12). Besides, female sex workers are also exposed to the risk of genital ulcers caused by infection with HSV-2. In recent years, an increasing prevalence of HSV-2 has been found around the world, particularly among high-risk groups such as female sex workers (13). Several clinical and epidemiological studies offered evidence for multiple interacttions between HIV and HSV-2 (14, 15).

In some meta-analysis and epidemiological studies, it has been demonstrated that HSV-2 can be regarded as a risk factor for HIV infection (16, 17). Furthermore, the status of antibodies to HSV-2 can be considered as a serological marker for the vulnerability to HIV infection in order to surveillance of infection controls (18, 19).

Considering the critical role of viral sexually transmitted infections and their complications and subsequences among female sex workers and their clients which pose a great burden on society and public health, it is worth investigating the prevalence of these infections in this high-risk group. Despite the numerous studies investigating the prevalence of STIs among female sex workers, there is no available study designed to characterize the global epidemiology of major viral STIs among this high-risk population. To the best of our knowledge, this is the first meta-analysis that aims to characterize the epidemiology of the three most important viral pathogens, including HBV, HCV, and HSV-2 among female sex workers around the world.

#### **Methods**

The protocol of this systematic review and meta-analysis was in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement guideline (20).

Search strategy: We carried out an electronic literature search to identify relevant publications on the prevalence of STIs including HBV, HCV, and HSV-2 infections in female sex workers in the world. Articles were identified through searches in electronic databases, including PubMed, Scopus, Google Scholar, Web of Science, and Embase from database inception to September 25, 2019.

Also, we manually searched relevant articles and bibliographic reference lists of the publiccations selected for inclusion in our metaanalysis.

All searched publications were imported to EndNote software version X8 (Thomson Reuters, California, USA) for further management.

Selection criteria: Studies were deemed eligible for inclusion if they reported: (1) original data published in peer-reviewed journals in the English language regarding the prevalence measure for each STI of interest among female sex workers; (2) the prevalence of hepatitis B surface antigen (HBsAg), anti-HCV antibody, anti-HSV-2 antibody in blood and saliva; (3) editorials, letters, commentaries, short reports, and English abstracts with sufficient data.

Author (Ref.)				n this systemat				
	Publication Study		Location Virus	Total	No.	No.	No.	
	Year	Period			sample	HBV	HCV	HSV-2
					size	positive	positive	positive
Frade (25)	2019	2015-2017	Brazil	HBV	153	3	-	-
Puga (26)	2018	2009-2010	Brazil	HCV	402	-	2	-
Ferreira-Júnior	2018	2016	Brazil	HBV/HCV	4245	16	38	-
(27)								
Julien (28)	2018	2017	Rwanda	HBV/HCV	3930	126	177	-
Zhu (29)	2017	2014	China	HSV-2	585	-	-	268
Zhu (30)	2017	2014-2015	China	HSV-2	1058	-	-	537
Vu (31)	2017	2013	Tanzania	HSV-2	1903	-	-	1095
Niama (32)	2017	2011-2012	Republic of	HBV/HCV	805	34	6	-
			Congo					
Mutagoma (33)	2017	2015	Rwanda	HBV/HCV	1978	50	28	-
de Matos (34)	2017	2009-2010	Brazil	HBV/HCV	402	6	3	-
Han (35)	2016	2012	China	HSV-2	1487	-	-	414
Yu (36)	2016	2010-2014	China	HCV	1026	-	27	-
Shen (37)	2016	2014	China	HCV	653	-	13	-
Adisesh (38)	2016	2010	India	HBV/HCV/	2532	58	53	72
				HSV-2				
Moayedi-Nia	2015	2012-2013	Iran	HBV/HCV	161	2	13	-
(39)								
Fan (40)	2015	2012-2013	China	HCV	622	-	44	-
Salyuk (41)	2015	2013	Ukraine	HCV	4806	-	841	-
Chen (42)	2015	2010-2012	China	HCV	51790	-	516	-
Luo (43)	2015	2012	China	HSV-2	833	-	-	570
Bugssa (44)	2015	2013	Ethiopia	HBV	319	19	-	-
Tang (45)	2014	2010	China	HCV	12622	-	128	-
Aho (46)	2014	2005-2006	Guinea	HSV-2	201	-	-	169
Kazerooni (47)	2014	2010-2011	Iran	HSV-2	278	-	-	27
Zhang (48)	2014	2011	China	HBV	600	74	-	-
Lwin (49)	2014	2009-2010	Myanmar	HCV	150	-	7	-
Li (50)	2014	NR	China	HSV-2	460	-	-	198
Fu (51)	2014	NR	China	HSV-2	575	-	-	176
Wang (52)	2014	2008-2012	China	HCV	759783	-	6021	-
Vandenhoudt	2013	2008	Kenya	HSV-2	479	-	-	404
(53)								
Zohrabyan (54)	2013	2009-2010	Moldova	HBV/HCV	658	60	149	_
Zhou (55)	2013	2010	China	HCV	12622	-	126	-
Valadez (56)	2013	2010	Libya	HBV/HCV	69	2	5	-
~	2013	2009	Brazil	HBV/HCV	134	5	12	-
Schuelter-								
Schuelter- Trevisol (57)	2012	2009	China	HSV-2	2453	-	-	1347
	2013			HBV/HCV	250	19	7	
Trevisol (57)	2013	2007	India	I DD V/DC V	230	12	/	-
Trevisol (57) Chen (58)		2007 2012	India Indonesia	HBV/HCV	200	8	1	-
Trevisol (57) Chen (58) Praseeda (59)	2013							
Trevisol (57) Chen (58) Praseeda (59) Kotaki (60)	2013 2013	2012	Indonesia	HBV/HCV	200	8	1	

### HBV, HCV, and HSV-2 infections in sex workers

				HSV-2				
Sarna (63)	2013	2010	India	HSV-2	529	-	_	321
Prestileo (64)	2013	1999-2008	Italy	HBV/HCV	239	7	4	-
Mollik (65)	2013	NR	Bangladesh	HBV	196	18		
Ghosh (66)	2012	NR	India	HBV/HSV-	45	0	-	13
Zermiani (67)	2012	1999-2007	Italy	HBV/HCV	345	12	3	-
Navadeh (68)	2012	2010	Iran	HSV-2	139	-	-	28
Wong (69)	2012	2005-2007	Hong Kong	HBV	503	43	-	-
Dunford (70)	2012	2008-2009	Vietnam	HBV	1000	99	-	-
Wang (71)	2012	2009	China	HSV-2	345	_	-	201
Dunford (72)	2012	2008-2009	Vietnam	HCV	1000	-	87	-
Cárcamo (73)	2012	2002-2003	Peru	HSV-2	4485	-	-	2718
Barua (74)	2012	2006	India	HCV	426	-	41	-
Delgado (75)	2011	2009	Nicaragua	HSV-2	825	-	-	641
Qyra (76)	2011	2008	Albania	HBV	92	7	-	-
Yang (77)	2011	2008	China	HSV-2	793	-	-	375
Kassak (78)	2011	2007-2008	Lebanon	HBV/HCV	103	0	0	-
Hemalatha (79)	2011	2005-2006	India	HSV-2	322	-	-	225
Kassaian (80)	2011	2009-2010	Iran	HBV/HCV	91	1	9	-
Braunstein (81)	2011	NR	Rwanda	HSV-2	800	-	-	478
Znazen (82)	2010	2007	Tunisia	HBV/HCV/ HSV-2	183	1	2	105
Todd (83)	2010	2006-2008	Afghanistan	HBV/HCV	520	34	10	-
Zermiani (84)	2010	1999-2008	Italy	HBV/HCV	395	12	3	-
Shethwala (85)	2009	2005-2006	India	HBV	300	10	-	-
Wang (86)	2009	2006	China	HSV-2	737	-	-	502
Cavlek (87)	2009	2003-2006	Croatia	HCV	272	-	11	-
Shahmanesh (88)	2009	2004-2005	India	HSV-2	326	-	-	182
Bautista (89)	2009	2000-2002	Argentina	HCV	625	-	26	-
Uusküla (90)	2008	2005-2006	Estonia	HCV	191	-	15	-
Yun (91)	2008	2003	South Korea	HSV-2	188	-	-	162
Ngo (92)	2008	2004	China	HCV/ HSV-2	310	-	24	140
Xu (93)	2008	2006	China	HSV-2	96			68
Forbi (94)	2008	2006-2007	Nigeria	HBV	720	123	-	-
Linhart (95)	2008	NR	Israel	HSV-2	300	-	-	180
Wang (13)	2008	2006	China	HSV-2	737	-	-	502
Taylor (96)	2008	1999	Scotland	HCV	98	-	63	-
Gul (97)	2008	2005	Turkey	HBV/HCV/ HSV-2	130	4	1	8 (IgM), 104 (IgG)
Barua (98)	2008	NR	India	HCV	426	_	41	(IgO) -
Monsalve-	2007	2004-2005	Venezuela	HCV	47	_	0	_
Castillo (99)	2007	2001 2003	, chezacia	110 7	1,			
Passos (100)	2007	NR	Brazil	HBV	449	3	-	_
Vandepitte (101)	2007	2002	Republic of Congo	HSV-2	501	-	-	293
Davies (102)	2007	1999-2000	Indonesia	HSV-2	165			149
Papadogeorgaki	2006	2005	Greece	HBV/HCV/	299	4	7	208

(103)				HSV-2				
Theng (104)	2006	2003-2004	Singapore	HSV-2	300	_	_	237
Pando (105)	2006	2000-2002	Argentina	HCV	602	-	26	-
Nigro (106)	2006	2001-2002	Italy	HBV/HCV	118	2	0	_
Dolar (107)	2006	NR	Turkey	HSV-2	483	-	-	290
Kweon (108)	2006	2004	South Korea	HBV/HCV	1527	102	21	-
Fox (109)	2006	1985-2000	UK	HSV-2	453	102		272
Cwikel (110)	2006	NR	Israel	HBV/HCV	43	3	4	-
O'Farrell (111)	2006	2002-2003	Vietnam	HSV-2	904	-	-	251
Didelot-	2006	2003-2004	Burkina	HSV-2	379	_	_	264
Rousseau (112)	2000	2003 200 1	Faso	115 , 2	3,7			201
Inciardi (113)	2006	2001-2003	USA	HCV	533	_	158	-
Chen (114)	2005	1999-2000	China	HSV-2	505	_	-	327
del Amo (115)	2005	2002	Spain	HBV/HCV	734	2	1	-
Nigam (116)	2005	NR	India	HBV	60	5	_	-
Nessa (117)	2004	2002	Bangladesh	HSV-2	400	-	_	138
Belza (118)	2004	1999-2000	Spain	HCV	579	_	5	-
Taketa (119)	2003	1996	Thailand	HBV/HCV	100	11	2	-
Uribe-Salas	2003	1998	Mexico	HSV-2	468	_	_	401
(120)	2003	1,,,0	Wienieo	115 V 2	100			101
Miyazaki (121)	2003	1999-2001	Japan	HBV/HCV	171	0	2	-
Qutub (122)	2003	NR	Bangladesh	HSV-2	463	_	_	438
Valdivia (123)	2003	NR	Peru	HCV	98	_	0	-
Risbud (124)	2003	1994-1999	India	HCV	933	_	13	-
Huo (125)	2003	2001-2002	Taiwan	HBV/HCV	1045	126	86	-
Camejo (126)	2003	1999	Venezuela	HBV/HCV	212	8	1	-
Bystrická (127)	2003	NR	Slovakia	HBV/HCV/	18	4	0	9
J				HSV-2				-
Uribe-Salas	2002	1998	Mexico	HBV	461	6	-	-
(128)								
Miyazaki (129)	2002	1999-2000	Japan	HBV/HCV	135	0	1	-
Othman (130)	2002	NR	Syria	HBV/HCV	102	11	2	-
Laurent (131)	2001	1988	Republic of	HCV	1144	-	76	-
			Congo					
Ishi (132)	2001	NR	Japan	HBV/HCV	308	2	10	-
Rahman (133)	2000	1998	Bangladesh	HSV-2	203	-	-	127
Limpakarnjanara	1999	1991-1994	Thailand	HSV-2	500	-	-	378
t(134)								
Ghebrekidan	1999	1995	Eritrea	HSV-2	107	-	-	86
(135)								
Conde-Glez	1999	1992	Mexico	HSV-2	997	-	-	606
(136)								
Suganuma (137)	1998	1996	Thailand	HCV	100	-	2	-
Juárez-Figueroa	1998	1992	Mexico	HBV	1498	3	-	-
(138)								
Dada (139)	1998	1990-1991	Nigeria	HSV-2	796			470
Ghebrekidan	1998	1995	Eritrea	HBV/HCV	107	3	3	-
(140)								
McDonnell (141)	1998	1991-1997	Ireland	HBV/HCV	99 (for	6	8	-
					HCV),			

					124 (for HBV)			
Uribe-Salas (142)	1997	1992	Mexico	HBV/HSV- 2	778 (for HBV), 757	3	-	493
					(For HSV-2)			
Tsakris (143)	1997	NR	Greece	HBV	230	8	-	-
Luksamijarulkul (144)	1997	1995	Thailand	HCV	200	-	19	-
Mesquita (145)	1997	1986-1990	Brazil	HBV/HCV	686	34	105	-
Sattar (146)	1996	NR	Bangladesh	HBV	164	16	-	-
Nakashima (147)	1996	1989-1992	Japan	HCV	604	-	61	-
Weber (148)	1995	1991-1993	Germany	HCV	364	-	5	-
Watts (149)	1994	1990	Somalia	HCV	236	-	4	-
Gotuzzo (150)	1994	1991-1992	Peru	HSV-2	399	-	-	328
Develouxl (151)	1994	1990	Niger	HCV	250	-	19	-
Wu (152)	1993	1988-1990	Taiwan	HCV	622	-	74	-
Ndumbe (153)	1993	1990-1992	Cameroon	HCV	104	-	16	-
Echevarria (154)	1993	1990	Spain	HCV	121	-	15	-
Tuncel (155)	1992	NR	Turkey	HBV	101	9	-	-
Orduna (156)	1992	NR	Spain	HBV/HCV	227	14	20	-
Gutierrez (157)	1992	NR	Spain	HCV	227	-	20	-
Wu (158)	1991	NR	Taiwan	HCV	223	-	23	-
Lee (159)	1991	1988-1999	Taiwan	HBV/HCV	196	42	14	-
Ramachandran (160)	1990	1988	Malaysia	HBV	98	6	-	-
Mak (161)	1990	1988-1989	Belgium	HBV/HCV	123	0	3	-
Hyams (162)	1990	1988	Mexico	HBV	354	3	-	-
Caballero (163)	1987	NR	Spain	HBV	80	8	-	-
Jama (164)	1987	1985-1986	Somalia	HBV	85	17	-	-
Goh (165)	1985	1982	Singapore	HBV	239	15	-	-
de Hoop (166)	1984	1982	Netherlands	HBV	128	3	-	-
Kaklamani (167)	1980	1978-1979	Greece	HBV	198	22	-	-
Frosner (168)	1975	1973	Germany	HBV	258	6	-	-

NR: Not reported; HBV: Hepatitis B virus; HCV: Hepatitis C virus; HSV-2: Herpes simplex virus 2.


Studies that met any one of the following criteria were excluded: (1) studies investigating the incidence of STIs among female sex workers; (2) publications about the incidence and the prevalence of STIs among transgenders, male sex workers, men who have sex with men, lesbians, clients of female sex workers (3) studies evaluating the prevalence of occult hepatitis B (OBI) among female sex workers. (4) Case reports, posters, review articles, and conference abstracts; (5) publica-

tions in languages other than English with non-English abstracts.

Data extraction and quality assessment: Two authors independently extracted data from eligible studies onto a pre-designed data extraction form using Microsoft Excel 2013 (Microsoft Corporation, Redmond, Washington, USA), and any disagreements were resolved by a third author.

Tunisia	1	0.55 (0.08-3.77)	NA, NA
Turkey	2	5.59 (1.94-15.05)	70.0%, P= 0.07
Venezuela	1	3.77 (1.90-7.36)	NA, NA
Vietnam	1	9.90 (8.20-11.91)	NA, NA

NA: Not applicable; ELISA: Enzyme-Linked Immunosorbent Assay. † Statistically significant



**Fig. 1.** Flowchart presenting the steps of literature search and selection

The following characters were extracted from each eligible study: the first author's last name, publication date, study date, study location, total sample size, type of sample, number of HBV, HCV, and HSV-2 -positive cases, diagnostic methods, and diagnostic indexes (for HSV-2).

A quality assessment of the retrieved studies was conducted using a modified checklist based on the guidelines of the strengthening the reporting of observational studies in epidemiology (STROBE). The checklist was comprised of 12 questions covering different

methodological perspectives. Studies were deemed eligible for the main meta-analysis if they achieved a validity score of at least 8 out of a maximum of 12. Data extraction was undertaken independently by two authors, and any discrepancies at this stage were resolved by discussion and consultation with a third author.

Statistical analysis: To estimate the pooled prevalence for each STI among female sex workers, a DerSimonian-Laird random-effects meta-analysis using the inverse variance method was conducted (21). The logit transformation was applied to stabilize the variance and normalize their distribution, and the Clopper-Pearson method was applied to calculate the 95% exact confidence intervals (CIs) for proportions (22). To identify the possible sources of heterogeneity, subgroup analyses were performed, based on the diagnostic method, study location, and diagnostic indexes. A time-trend graph was produced using GraphPad Prism 7.1 for Windows (GraphPad Software, La Jolla California USA). To measure the heterogeneity among the included publications, I-square statistics (I<sup>2</sup>) was carried out, in which the result is presented as a percentage. I<sup>2</sup> values of 25%, 50%, and 75% are indicative of low, moderate, and high levels of heterogeneity, respectively (23). All analyses were implemented in the R package Meta (24) (version 3.5.3 [2019-03-11], R Foundation for Statistical Computing, Vienna, Austria), and for all statistical tests, P values of less than 0.05 were considered statistically significant.

#### **Results**

**Literature search:** In the initial search, 2516 articles were identified from five electronic international databases. A total of 1146 duplicates was excluded, and then 1370 articles

were screened by title and abstract, which led to the elimination of 1057 articles.

The remaining 313 articles were checked for agreement with the inclusion and exclusion criteria by the full-text review.




Fig. 2. Forest plot of the prevalence of HBsAg among female sex workers.

After the full-text screening, 177 articles were excluded due to the following: 1) articles which were not relevant to the subject (n = 125), 2) articles which were determined the incidence of STIs (n = 19), 3) articles which were determined the risk factors of STIs (n = 19).

11), 4) articles with English abstracts with insufficient data (n = 6), 5) articles which did not differentiate between male and female sex workers (n=5), 6) articles in languages other than English with non-English abstracts (n=4), 7) review articles (n=2), 8) articles which had no full-text available (n=2), 9) conference abstract (n=1), 10) article which did not specify diagnostic marker of HBV (n=1), and 11) article which was determined the prevalence of occult HBV (OBI) among female sex workers.

Also, 10 relevant articles were found and included by a manual search of the reference lists of the identified articles. Based on the modified STROBE checklist, all papers were deemed to have good quality (obtained scores of 8 and above), and none of the papers was failed to reach score of 8. Finally, 146 articles were included in this systematic review and meta-analysis. Fig. 1 represents the process of literature retrieval and screening using a flow chart.

Study characteristics: The characteristics of eligible studies in this systematic review and meta-analysis are summarized in Table 1. One hundred forty-six studies with a total of 917816 female sex workers were included in this meta-analysis. The studies' publication dates ranged from 1975 to 2019, and they examined the population of 55 countries. The largest study included 759783 and the smallest included 18 female sex workers. Out of 146 studies included, 67 studies assessed HBsAg (in 34048 participants), 77 studies assessed anti-HCV antibodies (in 879725 participants), and 51 studies assessed anti-HSV-2 antibodies (in 34230 participants).

Most studies investigating the prevalence of HBsAg, anti-HCV antibody, and anti-HSV-2 antibodies were from Brazil (n=6), China (n=8), and China (n=14), respectively.

	Table 3. Subgroup a	nalysis of the preval	ence of HCV infection in fe	male sex workers	
Characteristics	Categories	No. of Studies	Pooled prevalence (%) (95% CI)	Heterogeneity test I <sup>2</sup> %, <i>p</i> -value	Differences between subgroups; χ² test (p-value)
Overall		77	3.39 (2.30-4.97)	99.5%, <i>P</i> <0.0001	<b>y</b> ,
Detection	CLEIA	2	1.00 (0.32-3.07)	0%, <i>P</i> =0.71	P=0.19
method	ELISA	68	3.69 (2.53-5.36)	99.3%, P<0.0001	
	Passive hemagglutination	2	2.71 (0.13-36.54)	89.4%, P<0.01	
G 1 4	Rapid Test	4	2.65 (0.37-16.52)	99.5%, <i>P</i> <0.01	D 0.011
Sample type	Blood	76	3.22 (2.19-4.71)	99.5%, P<0.0001	P< $0.01$ †
Ct., d., loootion	Saliva	1	64.29 (54.35-73.13)	NA, NA	D -0 014
Study location	Afghanistan	1	1.92 (1.04-3.54)	NA, NA	P< $0.01$ †
	Argentina	2	4.24 (3.24-5.52)	0%, <i>P</i> =0.89	
	Belgium	1	2.44 (0.79-7.29)	NA, NA	
	Brazil	5	2.34 (0.45-11.33)	98.5%, <i>P</i> <0.01	
	Cameroon	1	15.38 (9.64-23.65)	NA, NA	
	China	8	1.91 (1.33-2.72)	98.3%, <i>P</i> <0.01	
	Croatia	1	4.04 (2.25-7.15)	NA, NA	
	Eritrea	1	2.80 (0.91-8.33)	NA, NA	
	Estonia	1	7.85 (4.79-12.62)	NA, NA	
	Germany	1	1.37 (0.57-3.62)	NA, NA	
	Greece	1	2.34 (1.12-4.83)	NA, NA	
	India	5	3.88 (1.67-8.76)	96.1%, <i>P</i> <0.01	
	Indonesia	1	0.50 (0.07-3.46)	NA, NA	
	Iran	2	8.77 (5.84-12.96)	0%, P=0.62	
	Ireland	1	8.08 (4.09-15.33)	NA, NA	
	Israel	1	9.30 (3.54-22.30)	NA, NA	
	Italy	4	1.04 (0.57-1.89)	0%, P=0.63	
	Japan	4	2.96 (0.95-8.84)	88.5%, <i>P</i> <0.01	
	Lebanon	1	0.48 (0.03-7.21)	NA, NA	
	Libya	1	7.25 (3.05-16.26)	NA, NA	
	Moldova	1	22.64 (19.61-26.00)	NA, NA	
	Myanmar	1	4.67 (2.24-9.46)	NA, NA	
	Niger	1	7.60 (4.90-11.61)	NA, NA	
	Panama	1	0.20 (0.05-0.80)	NA, NA	
	Peru	1	0.51 (0.03-7.56)	NA, NA	
	Republic of Congo	2	2.34 (0.26-17.80)	96.4%, <i>P</i> <0.01	
	Republic of Mauritius	1	47.46 (41.82-53.16)	NA, NA	
	Rwanda	2	2.57 (0.82-7.79)	97.0%, <i>P</i> <0.01	
	Scotland		64.29 (54.35-73.13)	97.0%, P<0.01 NA, NA	
		1	, , , , , , , , , , , , , , , , , , , ,		
	Slovakia	1	2.63 (0.16-30.96)	NA, NA	
	Somalia	1	1.69 (0.64-4.43)	NA, NA	
	South Korea	1	1.38 (0.90-2.10)	NA, NA	
	Spain	5	3.70 (1.41-9.35)	91.4%, <i>P</i> <0.01	
	Syria	1	1.96 (0.49-7.50)	NA, NA	
	Taiwan	4	9.49 (7.50-11.95)	60.0%, <i>P</i> =0.06	
	Thailand	3	3.89 (1.11-12.79)	76.6%, <i>P</i> =0.01	

NA, NA


	Turkey	1	0.77 (0.11-5.25)	NA, NA	
	Ukraine	1	17.50 (16.45-18.60)	NA, NA	
	USA	1	29.64 (25.92-33.66)	NA, NA	
	Venezuela	2	0.61 (0.12-2.99)	0%, <i>P</i> =0.65	
	Vietnam	1	8.70 (7.10-10.61)	NA, NA	
CLEIA: Che	emiluminescence enzyme ir	nmunoassay; ELISA	: Enzyme-Linked Immunos	sorbent Assay; NA: No	t applicable.
† Statisticall	y significant				

1.09 (0.27-4.26)

Tunisia

	Table 4. Subgroup an	alysis of the serop	prevalence of HSV-2 in	female sex workers	
		-			
Characteristics	Categories	No. of Studies	Pooled prevalence (%) (95% CI)	Heterogeneity test I <sup>2</sup> %, p-value	Differences between subgroups; χ² test (p-value)
Overall		51	60.48 (55.43-56.31)	98.6%, <i>P</i> <0.0001	<u> </u>
Detection	ELISA	43	57.47 (51.71-63.03)	98.7%, P<0.0001	P=0.02†
method	Immunoelectrotransference	1	65.13 (61.66-68.44)	NA, NA	
	Rapid Test	2	72.54 (41.06-90.92)	97.9%, <i>P</i> <0.01	
	Western blotting	4	77.13 (63.82-86.58)	97.6%, <i>P</i> <0.01	
Detection	IgM antibody	5	10.31 (3.88-24.66)	96.7%, P<0.01	P<0.01†
index	IgG antibody	47	64.84 (60.73-68.75)	98.1%, <i>P</i> <0.0001	
Study location	Bangladesh	3	71.19 (29.00-93.73)	99.2%, <i>P</i> <0.01	P<0.01†
	Burkina Faso	1	69.66 (64.84-74.08)	NA, NA	·
	China	14	53.10 (45.54-60.51)	98.3%, <i>P</i> <0.01	
	Eritrea	1	80.37 (71.76-86.84)	NA, NA	
	Greece	1	69.57 (64.11-74.52)	NA, NA	
	Guinea	1	84.08 (78.35-88.51)	NA, NA	
	India	5	35.84 (9.82-74.13)	99.6%, <i>P</i> <0.01	
	Indonesia	1	90.30 (84.76-93.97)	NA, NA	
	Iran	2	14.12 (6.66-27.48)	88.2%, P<0.01	
	Israel	1	60.00 (54.35-65.40)	NA, NA	
	Kenya	1	84.34 (80.81-87.33)	NA, NA	
	Mexico	3	71.90 (57.67-82.78)	97.6%, <i>P</i> <0.01	
	Nicaragua	1	77.70 (74.73-80.41)	NA, NA	
	Nigeria	1	59.05 (55.59-62.41)	NA, NA	
	Panama	1	74.17 (71.37-76.79)	NA, NA	
	Peru	2	72.57 (47.39-88.60)	98.5%, <i>P</i> <0.01	
	Republic of Congo	1	58.48 (54.11-62.72)	NA, NA	
f 07-01-	Rwanda	1	59.75 (56.31-63.10)	NA, NA	
	Singapore	1	79.00 (74.02-83.24)	NA, NA	
	Slovakia	1	50.00 (28.42-71.58)	NA, NA	
	South Korea	1	86.17 (80.46-90.41)	NA, NA	
<b>∃</b> ⊇)	Tanzania	1	57.54 (55.31-59.74)	NA, NA	
	Thailand	1	75.60 (71.64-79.17)	NA, NA	
41 m	Tunisia	1	57.38 (50.11-64.34)	NA, NA	
	Turkey	2	74.99 (42.70-92.34)	96.4%, <i>P</i> <0.01	
<b>T</b>	UK	1	60.04 (55.46-64.46)	NA, NA	
Ĕ	Vietnam	1	27.77 (24.94-30.78)	NA, NA	

ELISA: Enzyme-Linked Immunosorbent Assay; NA: Not applicable; † Statistically significant.



**Fig. 3.** Forest plot of the prevalence of HCV infection among female sex workers.


**Prevalence of HBsAg marker among female sex workers:** The pooled prevalence of HBsAg was determined amongst 34048 female sex workers from 41 countries in the world. The overall prevalence of HBsAg was 3.76% (95% CI: 2.99%-4.72%), and the details are presented in a forest plot (Fig. 2). However, there was significant heterogeneity between the studies ( $I^2 = 93.2\%$ , P < 0.01), and the studies

met the random-effects model. Table 2 shows subgroup analyses of the prevalence of HBsAg in female sex workers stratified by detection method and study location. Subgroup analysis according to study location showed that the lowest pooled prevalence of HBsAg was seen in the Republic of Mauritius (0.17%; 95% CI: 0.01%-2.64%), while the highest pooled prevalence was found in Slovakia (22.22%; 95% CI: 8.60%-46.47%).

The majority of studies on the prevalence of HBsAg in female sex workers was conducted in Brazil (6 studies), followed by India (5 studies), Italy (4 studies), and Mexico (4 studies). According to the subgroup analysis results for detection method, the pooled prevalence of HBsAg was higher using the radioimmunoassay (9.93%; 95% CI: 5.97%-16.07%) than using the latex agglutination (8.33%; 95% CI: 3.51%-18.51%), ELISA (3.39%; 95% CI: 2.62%-4.38%), passive hemagglutination (3.16%; 95% CI: 1.86%-5.33%), and Rapid Test (1.89%; 95% CI: 0.4%-7.27%), the differences were statis-tically significant (P<0.01). In all studies, blood samples were used to evaluate the presence of HBsAg, and thus, there is no subgroup analysis of sample type.

**Prevalence of HCV infection among female sex workers**: The pooled prevalence of HCV infection was calculated amongst 879725 female sex workers from 42 countries across the world. The overall prevalence of HCV infection was 3.39% (95% CI: 2.30%-4.97%), and outputs of meta-analyses are depicted in a forest plot (Fig. 3). However, a high degree of heterogeneity existed among the included studies ( $I^2 = 99.5\%$ , P < 0.0001), and therefore, the studies met the random-effects model. Table 3 presents subgroup analyses of the prevalence of HCV infection among female sex workers stratified by detection method, sample type, and study location.

In the subgroup analysis according to study location, the highest prevalence was observed in Scotland (64.29%; 95% CI: 54.35%-73.13%), whereas the lowest prevalence was found in Panama (0.20%; 95% CI: 0.05%-0.80%).



**Fig. 4.** Forest plot of the seroprevalence of HSV-2 among female sex workers.

Most of the studies investigating the prevalence of HCV infection among female sex workers were performed in China (8 studies), followed by Brazil (5 studies), India (5 studies), and Spain (5 studies). Based on the subgroup analysis results for detection method, more HCV positive cases were detected by ELISA (3.69%; 95% CI: 2.53%-5.36%) than passive hemagglutination (2.71%; 95% CI: 0.13%-36.54%), Rapid Test (2.65%; 95% CI: 0.37%-16.52%), chemiluminescence enzyme immunoassay (CLEIA) (1.00%; 95% CI: 0.32%-3.07%). However, the differences were not statistically significant (*P*=0.19).

Furthermore, our subgroup analysis of sample type revealed that the prevalence of HCV infection was higher using the saliva samples (64.29%; 95% CI: 54.35%-73.13%) compared to the blood samples (3.22%; 95% CI: 2.19%-

4.71%), and the difference was statistically significant (P<0.01).

Prevalence of HSV-2 infection among female sex workers: A random-effects metaanalysis was performed to estimate the pooled seroprevalence of HSV-2 in 34230 female sex workers from 27 countries throughout the world. The pooled seroprevalence of HSV-2 was 60.48% (95% CI: 55.43%–65.31%). Totally, 47 studies (in 31236 participants) were conducted on the seroprevalence of IgG antibody, and 5 studies (in 3124 participants) were assessed the seroprevalence of IgM antibody. The overall prevalence of HSV-2 and IgM antibodies worldwide was IgG 64.84% (95% CI: 60.73%–68.75%) 10.31% (95% CI: 3.88%–24.66%), respectively, and the results of the meta-analysis are visually illustrated in a forest plot in Fig. 4. Table 4 displays subgroup analyses of the seroprevalence of HSV-2 among female sex workers, based on antibody class and study location. The seroprevalence of HSV-2 ranged from 2.84%-28.89% for IgM antibody and 27.77%-94.60% for IgG antibody.

In the subgroup analysis according to study location, the highest seroprevalence of HSV-2 was observed in Indonesia (90.30%; 95% CI: 84.76%-93.97%), whereas the lowest seroprevalence was found in Iran (14.12%; 95% CI: 6.66%-27.48%), respectively.

**Time trend analysis**: Time trend analysis was performed to investigate changes in the prevalence of STIs of interest over time in the world (Fig. 5). According to this analysis, the prevalence of HBsAg was the highest (11.11%; 95% CI: 7.43%-16.30%) between the years 1975 and 1979. From 2010 until 2019, the number of HBsAg-positive cases among female sex workers was decreased, and the lowest prevalence was seen between the years of 2015 and 2019 (1.58%; 95% CI: 0.69%-3.60%). Regarding HCV infection, the prevalence was high between the years of 1985 and 1994 (~8%). However, a dramatic decrease was observed during the period 1995-1999 (3.4%; 95% CI: 0.48%-20.51%), and from 2005 to 2019, a decreasing trend was observed in HCV prevalence among female sex workers.




Fig. 5. Time trends in the prevalence of each STI of interest among female sex workers.

Our results have shown that similar to HBsAg, the prevalence of HCV infection was the lowest between the years of 2015 and 2019 (1.81%; 95% CI: 0.59%-5.42%).

Furthermore, we found there was no remarkable change in the seroprevalence of HSV-2 among female prostitutes during the period 1990-2009. Nonetheless, a decrease in the seroprevalence was observed between the years 2010-2014.

#### **Discussion**

sex workers or prostitutes considered as one of the high-risk groups for sexually transmitted diseases and transmission of STIs and blood-borne infections like HBV and HCV. The large majority of prostitutes has been involved in unsafe sexual practice, generally owing to the lack of skills of condom use by their sexual partners. The prevalence estimates of HBsAg in female sex workers calculated in our current meta-analysis indicate that in most countries, the number of HBsAgpositive cases is significantly higher than those observed in the general population. For instance, our meta-analysis indicated that the pooled prevalences of HBsAg among Chinese and Thai female prostitutes were 12.3% and

11%, respectively, whereas according to the results of the previous meta-analysis, the estimates in the general population were 6.8% and 5.1%, respectively (169, 170).

According to the results of our meta-analysis, the highest prevalence rates of HBV infection were mainly observed among female sex workers in the Asia-Pacific and sub-Saharan Africa regions. In comparison with other countries, China (12.3%), Taiwan (16%), Thailand (11%), Hong Kong (8.5%), Malaysia (6.1%), Singapore (6.2%), Bangladesh (9.4%), Vietnam (10%) (From Asia-Pacific region), Somalia (20%) and Nigeria (17%) (From sub-Saharan Africa region) had a higher prevalence of HBV infection among female sex workers. This arises from the fact that HBV is endemic in many parts of the Asia-Pacific and sub-Saharan Africa regions (171).

Similarly, our meta-analysis also revealed the HCV infection occur at a higher rate among female sex workers than the general population. For example, according to our meta-analysis results, the pooled prevalence of HCV infection among Iranian, Chinese, and Cameroon female sex workers was 8.7%, 1.9%, and 15.3%, respectively, while the corresponding values in the general population calculated in

previous meta-analyses were 0.3% (172), 0.9% (173), and 3.6% (174), respectively.

Regarding the aforementioned results, it can be confirmed that female sex workers are at increased risk of both HBV and HCV acquisition.

Some countries such as Belgium, Japan, Mexico, and Brazil, have lower prevalences of HBV and HCV infections in female sex workers compared to the other countries. The explanation for these low prevalence estimates can be attributed to some influential factors such as prostitutes' knowledge and awareness about STDs transmission and prevention modes, regular STD screening by rapid qualitative tests, using barrier methods such as condoms, and HBV vaccination of sex workers. For example, the Ministry of Health of Brazil has been developed campaigns to distribute free condoms at health centers (26). Nevertheless, condom failure (breakage or slippage) is a common problem among female sex workers, especially those who have experienced client-perpetuated violence.

Condom breakage is estimated to be 0.8% to 40.7%, and condom slippage during sexual intercourse is estimated to be 13.1% and 19.3% (175).

The risk of STD infection among female sex workers is significantly related to the type of sex work environments. The current metaanalysis showed that the pooled prevalences of HBV and HCV infection among Japanese prostitutes are low. In Japan, the most common workplace for female sex workers is massage parlors with cell baths (121). Street-based female sex workers are more likely than indoor-based sex workers to experience poverty, poor health, and social exclusion (176). A similar situation has been reported for Mexican female sex workers so that women working at street sites had higher rates of STIs such as HSV-2 and HBV compared to women working in bars and massage parlors (142). It should be noted that due to anal tissue is more prone to tears, the practice of anal sex with female sex workers is associated with an increased risk of STIs transmission (177, 178). Since that anal sex was an infrequent practice among Japanese and Mexican female sex

workers included in this meta-analysis (121, 129, 138), it might justify lower prevalences of HBV and HCV among female sex workers in these countries. Another explanation for the low prevalence of viral hepatitis in Mexican prostitutes may partly be due to the absence of injecting drug use (138). Injection drug use is an important public health concern and a major risk factor for increasing the prevalence of STIs among female prostitutes (34).

The overall pooled seroprevalence of HSV-2 was 60.5% among female sex workers, indicating a high exposure of the population to the infection. The seroprevalence of HSV-2 ranged from 14.1% in Iran to 90.3% in Indonesia. HSV-2 infection is one of the most common STIs worldwide and is a major causative agent of genital ulcers (58). The virus can infect anogenital regions and creates a potential and lifelong source of viral transmission to sexual partners (179). In previous studies, strong multiple interactions observed between HSV-2 and HIV, and also HSV-2 recognized as a risk factor for HIV infection. For example, in the meta-analysis carried out by Freeman et al., infection with HSV 2 was associated with a three times increase in the risk of acquiring HIV in the general population (180). The results presented by Schacker et al. also confirmed an increased efficiency of the sexual HIV transmission with genital herpes infection (181). Furthermore, the HSV-2 antibody status is often used as a serological marker of HIV susceptibility (58). Hence, routine screening of HSV-2 antibody status can be used as a predictive factor of high-risk behaviors in female sex workers and will be helpful to develop efficient intervention and surveillance programs.

The present study has some limitations; (1) there were no data regarding the prevalence of HBV, HCV, and HSV-2 among female sex workers especially for countries in Northern America and Europe. We did not find any published articles for countries such as Canada, Cuba, Haiti, and El Salvador from Northern America, and France, Sweden, Norway, Poland, Romania, Portugal, Russia, and many other countries from Europe; (2) the number of studies was limited in countries such as USA,

UK, and Germany, with only one study included; (3) In more conservative settings and Islamic countries such as Pakistan, Iran, Saudi Arabia, UAE, and Afghanistan, female sex workers have extensively remained as a hidden population. Most female sex workers in these countries do not attend public STD clinics, and therefore, estimation of the prevalence of the STIs of interest is not precise in our metaanalysis. (4) Included studies in this metaanalysis have used different diagnostic kits for measuring the prevalence of viral infections. It should be noted that the sensitivity and specificity of each kit are varied, and therefore. the difference between results can be attributed partly to this feature.

#### Conclusion

This meta-analysis concluded that the prevalence of STIs including HBV, HCV, and HSV-2 infections among female sex workers is considerably high, particularly in some regions, and highlights the importance of STIs screening of this high-risk, vulnerable, and neglected group to public health.

It is required for female sex workers to implement comprehensive health promotion programs, develop inexpensive, sensitive, and rapid tests, improve prevention programs such as vaccination (for HBV), and using barrier methods like condoms.

## Acknowledgment

Not applicable.

#### **Conflict of interest**

The authors have no conflict of interest.

## **Funding**

This study was not financially supported by any individual, agency, or institution.

#### References

- 1. Bellhouse C, Crebbin S, Fairley CK, Bilardi JE. The impact of sex work on women's personal romantic relationships and the mental separation of their work and personal lives: A mixed-methods study. PloS One. 2015; 10(10):e0141575.
- 2. Sawicki DA, Meffert BN, Read K, Heinz AJ. Culturally competent health care for sex workers: an examination of myths that stigmatize sex work and hinder access to care. Sex Relation Ther. 2019:1-17.
- 3. Mamulwar M, Godbole S, Bembalkar S, Kamble P, Dulhani N, Yadav R, et al. Differing HIV vulnerability among female sex workers in a high HIV burden Indian state. PloS One. 2018;13(2):e0192130.
- 4. Szwarcwald CL, Damacena GN, de Souza-Júnior PRB, Guimarães MDC, de Almeida WdS, de Souza Ferreira AP, et al. Factors associated with HIV infection among female sex workers in Brazil. Medicine. 2018; 97(1 Suppl).
- 5. Hernandez I, Reina-Ortiz M, K. Yusuf K, Ochoa T, Rosas C, Sharma V, et al. Risk factors for HIV and STI among female sex workers in a high HIV prevalent region of Ecuador. Cogent Med. 2019;6(1):1565292.
- 6. Carmona-Gutierrez D, Kainz K, Madeo F. Sexually transmitted infections: old foes on the rise. Microbial Cell 2016;3(9):361.
- 7. Apidechkul T. Sexual behaviors and seroprevalence of HIV, HBV, and HCV among hill tribe youths of Northern Thailand. 2019;19(1):1-17.
- 8. https://www.who.int/news-room/fact-sheets/detail/hiv-aids.
- 9. https://www.who.int/news-room/fact-sheets/detail/hepatitis-b.
- 10. https://www.who.int/news-room/fact-sheets/detail/hepatitis-c.
- 11. Sadeghi F, Bokharaei-Salim F, Salehi-Vaziri M, Monavari SH, Alavian SM, Salimi S, et al. Associations between human TRIM22 gene expression and the response to combination therapy with Peg-IFN $\alpha$ -2a and ribavirin in Iranian patients with chronic hepatitis C. J Med Virol. 2014;86(9):1499-506.
- 12. Bokharaei-Salim F, Keyvani H, Monavari SH, Alavian SM, Fakhim S, Nasseri S. Distribution of hepatitis C virus genotypes among azerbaijani patients in capital city of iran-tehran. Hepat Mon. 2013;13(9): e13699
- 13. Wang H, Wang N, Chen RY, Sharp GB, Ma Y, Wang G, et al. Prevalence and predictors of herpes simplex virus type 2 infection among female sex workers in Yunnan Province, China. Int J STD & AIDS. 2008;19(9):635-9.
- 14. Zhu J, Hladik F, Woodward A, Klock A, Peng T, Johnston C, et al. Persistence of HIV-1 receptor—positive cells after HSV-2 reactivation is a potential mechanism for increased HIV-1 acquisition. Nat Med. 2009;15(8): 886.
- 15. Corey L. Synergistic copathogens—HIV-1 and HSV-2. N Engl J Med. 2007;356(8):845-6.
- 16. Looker KJ, Elmes JA, Gottlieb SL, Schiffer JT, Vickerman P, Turner KM, et al. Effect of HSV-2 infection on subsequent HIV acquisition: an updated

- systematic review and meta-analysis. Lancet Infect Dis. 2017;17(12):1303-16.
- 17. Wald A, Link K. Risk of human immunodeficiency virus infection in herpes simplex virus type 2–sero-positive persons: a meta-analysis. J Infect Dis. 2002;185 (1):45-52.
- 18. Wang H, Reilly K, Smith M, Brown K, Jin X, Xu J, et al. Herpes simplex virus type 2 incidence and associated risk factors among female sex workers in a high HIV-prevalence area of China. Int J STD AIDS. 2013;24(6):441-6.
- 19. Cowan FM, Johnson AM, Ashley R, Corey L, Mindel A. Antibody to herpes simplex virus type 2 as serological marker of sexual lifestyle in populations. BMJ. 1994;309(6965):1325-9.
- 20. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. Ann Intern Med. 2009;151(4):264-9.
- 21. DerSimonian R, Laird N. Meta-analysis in clinical trials. Control Clin Trials.1986;7(3):177-88.
- 22. Newcombe RG. Two-sided confidence intervals for the single proportion: comparison of seven methods. Stat Med.1998;17(8):857-72.
- 23. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. BMJ. 2003; 327(7414):557-60.
- 24. Schwarzer G. meta: An R package for meta-analysis. R News. 2007;7(3):40-5.
- 25. Frade PC, Raiol NC, da Costa LM, Pinheiro LM, Silva-Oliveira GC, Pinho JR, et al. Prevalence and genotyping of hepatitis B virus: a cross-sectional study conducted with female sex workers in the Marajó Archipelago, Brazil. Int J STD & AIDS. 2019:30(9): 902-910.
- 26. Puga MAM, Bandeira LM, Weis SMdS, Fernandes FRP, Castro LS, Tanaka TSO, et al. High-risk behaviors for hepatitis B and C infections among female sex workers. Rev Soc Bras Med Trop. 2018;51(2):198-202.
- 27. da Costa Ferreira-Júnior O, Guimarães MDC, Damacena GN, de Almeida WdS, de Souza-Júnior PRB, Szwarcwald CL. Prevalence estimates of HIV, syphilis, hepatitis B and C among female sex workers (FSW) in Brazil, 2016. Medicine. 2018;97(1 Suppl).
- 28. Mutagisha JN, Ingabire R, Unyuzimana MA, Sinabamenye R, Bizimana J, Hoagland AB, et al., editors. Hepatitis B and C Prevalence Among Female Sex Workers in Kigali, Rwanda. AIDS RESEARCH AND HUMAN RETROVIRUSES; 2018: MARY ANN LIEBERT, INC 140 HUGUENOT STREET, 3RD FL, NEW ROCHELLE, NY 10801 USA.
- 29. Zhu J, Yuan R, Hu D, Zhu Z, Wang N, Wang B. HIV prevalence and correlated factors of female sex workers and male clients in a border region of Yunnan Province, China. Int J STD & AIDS 2018;29(5):424-34. 30. Zhu J, Yuan R, Hu D, Zhu Z, Yang X, Wang N, et al. Consecutive cross-sectional survey of prevalence of HIV infection/STD and related factors in Vietnamese female sex workers at a China-Vietnam border area,

- 2014-2015. Zhonghua liu xing bing xue za zhi= Zhonghua liuxingbingxue zazhi 2017;38(5):638-42.
- 31. Vu L, Misra K. High burden of HIV, syphilis and HSV-2 and factors associated with HIV infection among female sex workers in Tanzania: Implications for early treatment of HIV and pre-exposure prophylaxis (PrEP). AIDS Behav. 2018;22(4):1113-21.
- 32. Niama FR, Bongolo NCL, Mayengue PI, Mboussou FF, Bayonne ESK, Nzingoula FMK, et al. A study on HIV, Syphilis, and Hepatitis B and C virus infections among female sex workers in the Republic of Congo. Arch Public Health. 2017;75(1):21.
- 33. Mutagoma M, Nyirazinyoye L, Sebuhoro D, Riedel DJ, Ntaganira J. Syphilis and HIV prevalence and associated factors to their co-infection, hepatitis B and hepatitis C viruses prevalence among female sex workers in Rwanda. BMC Infect Dis. 2017;17(1):525.
- 34. Matos MAd, França DDdS, Carneiro MAdS, Martins RMB, Kerr LRFS, Caetano KAA, et al. Viral hepatitis in female sex workers using the Respondent-Driven Sampling. Rev Saude Publica. 2017;51:65.
- 35. Han L, Zhou C, Li Z, Poon AN, Rou K, Fuller S, et al. Differences in risk behaviours and HIV/STI prevalence between low-fee and medium-fee female sex workers in three provinces in China. Sex Transm Infect. 2016;92(4):309-15.
- 36. Yu YJ, Li X, Tam CC, Zhou Y, Chen Y, Shen Z. Demographic and behavioral correlates of HIV/STI among Vietnamese female sex workers in southwest China. AIDS care. 2016;28(11):1455-60.
- 37. Shen H, Zou H, Huang S, Liu F, Zhao P, Chen L, et al. Depression and HIV risk behaviors among female sex workers in Guangdong, China: a multicenter cross-sectional study. BioMed Res Int 2016;2016:6986173.
- 38. Adisesh M, Natesan T, Elumalai S, Subramanian K. A seroprevalence evaluation study of hepatitis B virus, hepatitis C virus and herpes simplex virus-2 among a special study group. Int J Med Res & Health Sci. 2016; 5(5):196-9.
- 39. Moayedi-Nia S, Bayat Jozani Z, Esmaeeli Djavid G, Entekhabi F, Bayanolhagh S, Saatian M, et al. HIV, HCV, HBV, HSV, and syphilis prevalence among female sex workers in Tehran, Iran, by using respondent-driven sampling. Aids Care. 2016;28(4):487-90.
- 40. Fan Y-G, Liu J-J, Zhang Y-J, Dai S-Y, Li M-Q, Ye D-Q. HIV, other sexually transmitted infections, and risk behaviors among female sex workers in Liuzhou, China. Int J Gynaecol Obstet. 2015;128(1):18-22.
- 41. Salyuk T, Sazonova I. Factors associated with hepatitis C virus (HCV) among female sex workers (FSWs) in Ukraine. J Clin Virol. 2015;69:245.
- 42. Chen Y, Shen Z, Morano JP, Khoshnood K, Wu Z, Lan G, et al. Bridging the epidemic: a comprehensive analysis of prevalence and correlates of HIV, hepatitis C, and syphilis, and infection among female sex workers in Guangxi Province, China. PloS One. 2015;10(2): e0115311.
- 43. Luo L, Xu J-J, Wang G-X, Ding G-W, Wang N, Wang H-B. Vaginal douching and association with sexually transmitted infections among female sex

- workers in a prefecture of Yunnan Province, China. Int J STD & AIDS. 2016;27(7):560-7.
- 44. Bugssa G, Dessalegn B, Dimtsu B, Berhane Y. Prevalence and factors associated with HIV and hepatitis B virus infections among female commercial sex workers in Mekelle, Ethiopia: cross sectional study. Int J Pharm Sci Res. 2015;6(1):135.
- 45. Tang Z, Zhang C, Li X, Liu Y, Su S, Zhou Y, et al. HIV risk among female sex workers with different patterns of drug use behaviors in Southwest China: a cross-sectional study. AIDS care. 2015;27(3):293-300.
- 46. Aho J, Koushik A, Coutlée F, Diakité SL, Rashed S. Prevalence of HIV, human papillomavirus type 16 and herpes simplex virus type 2 among female sex workers in Guinea and associated factors. Int J STD & AIDS. 2014;25(4):280-8.
- 47. Kazerooni PA, Motazedian N, Motamedifar M, Sayadi M, Sabet M, Lari MA, et al. The prevalence of human immunodeficiency virus and sexually transmitted infections among female sex workers in Shiraz, South of Iran: by respondent-driven sampling. Int J STD & AIDS. 2014;25(2):155-61.
- 48. Zhang T, Yang Y, Yu F, Zhao Y, Lin F, Minhas V, et al. Kaposis sarcoma associated herpesvirus infection among female sex workers and general population women in Shanghai, China: a cross-sectional study. BMC Infect Dis. 2014;14(1):58.
- 49. Lwin AA, Aung MM, Hpoo H, Aung PP, Tun WM, Oo KM, et al. Seroprevalence of hepatitis C virus infection among female commercial sex workers attending the Sexually Transmitted Diseases Clinic, Yangon General Hospital. Hepatol Int. [Conference Abstract]. 2014;8(1):S209.
- 50. Li Z, Li D, Liu H, Jiang Z, Zhu L, Wang N. Study on the prevalence and associated risk factors related to HIV, syphilis, herpes simplex virus-2 among female sex workers in Jiaozhou, Shandong province. Zhonghua Liu Xing Bing Xue Za Zhi. 2014;35(10):1099-104.
- 51. Fu X, Wang B, Li Y, Yang C, Lin G, Lin P. Prevalence of HIV and sexually transmitted diseases as well as related associated risk factors among middle/low level female sex workers in a city in Guangdong province. Zhonghua Liu Xing Bing Xue Za Zhi. 2014;35 (5):510-3.
- 52. Wang L, Tang W, Wang L, Qian S, Li Y-g, Xing J, et al. The HIV, syphilis, and HCV epidemics among female sex workers in china: results from a serial cross-sectional study between 2008 and 2012. Clin Infect Dis. 2014;59(1):e1-e9.
- 53. Vandenhoudt HM, Langat L, Menten J, Odongo F, Oswago S, Luttah G, et al. Prevalence of HIV and other sexually transmitted infections among female sex workers in Kisumu, western Kenya, 1997 and 2008. PloS One. 2013;8(1):e54953.
- 54. Zohrabyan L, Johnston LG, Scutelniciuc O, Iovita A, Todirascu L, Costin T, et al. Determinants of HIV infection among female sex workers in two cities in the Republic of Moldova: the role of injection drug use and sexual risk. AIDS Behav. 2013;17(8):2588-96.

- 55. Zhou Y, Li X, Zhang C, Tan G, Stanton B, Zhang X, et al. Rates of HIV, syphilis, and HCV infections among different demographic groups of female sex workers in Guangxi China: Evidence from 2010 national sentinel surveillance data. AIDS Care. 2013;25(11):1433-41.
- 56. Valadez JJ, Berendes S, Jeffery C, Thomson J, Othman HB, Danon L, et al. Filling the knowledge gap: measuring HIV prevalence and risk factors among men who have sex with men and female sex workers in Tripoli, Libya. PloS One. 2013;8(6):e66701.
- 57. Schuelter-Trevisol F, Custodio G, Silva ACBd, Oliveira MBd, Wolfart A, Trevisol DJ. HIV, hepatitis B and C, and syphilis prevalence and coinfection among sex workers in Southern Brazil. Rev Soc Bras Med Trop. 2013;46(4):493-7.
- 58. Chen S, Yin Y, Chen X, Wang H, Yu Y, Wei W, et al. Seropositivity and Risk Factors for Herpes Simplex Virus Type 2 Infection among Female Sex Workers in Guangxi, China. PloS one. 2013;8(7):e69697.
- 59. Desai Praseeda S, Anuradha D. A Study on the HBV and the HCV Infections in Female Sex Workers and their Co-Infection with HIV. J Clin Diag Res. 2013;7(2): 234.
- 60. Kotaki T, Khairunisa SQ, Sukartiningrum SD, Arfijanto MV, Utsumi T, Normalina I, et al. High prevalence of HIV-1 CRF01\_AE viruses among female commercial sex workers residing in Surabaya, Indonesia. PloS One. 2013;8(12):e82645.
- 61. Johnston LG, Corceal S. Unexpectedly high injection drug use, HIV and hepatitis C prevalence among female sex workers in the Republic of Mauritius. AIDS Behav. 2013;17(2):574-84.
- 62. Hakre S, Arteaga G, Núñez AE, Bautista CT, Bolen A, Villarroel M, et al. Prevalence of HIV and other sexually transmitted infections and factors associated with syphilis among female sex workers in Panama. Sex Transm Infect. 2013;89(2):156-64.
- 63. Sarna A, Friedland BA, Srikrishnan AK, Katzen LL, Tun W, Abbott SA, et al. Sexually transmitted infections and reproductive health morbidity in a cohort of female sex workers screened for a microbicide feasibility study in Nellore, India. Glob J Health Sci. 2013;5(3):139.
- 64. Prestileo T, Orlando G, Di FL, Dalle EN, Cassarà G, Bellipanni P, et al. Sexually transmitted infections in a cohort of female sex workers in Palermo. Recenti Prog Med. 2013;104(12):615-8.
- 65. Mollik AH. Investigations on human immunodeficiency virus and hepatitis B virus infections among the sex workers of Daulatdia-Ghat in Rajbari district Bangladesh. Hepatol Int. [Conference Abstract]. 2013;7:S161.
- 66. Ghosh I, Ghosh P, Bharti AC, Mandal R, Biswas J, Basu P. Prevalence of human papillomavirus and coexistent sexually transmitted infections among female sex workers, men having sex with men and injectable drug abusers from eastern India. Asian Pac J Cancer Prev. 2012;13(3):799-802.
- 67. Zermiani M, Mengoli C, Rimondo C, Galvan U, Cruciani M, Serpelloni G. Prevalence of sexually transmitted diseases and hepatitis C in a survey of

- female sex workers in the north-East of Italy. Open AIDS J. 2012;6:60.
- 68. Navadeh S, Mirzazadeh A, Mousavi L, Haghdoost A, Fahimfar N, Sedaghat A. HIV, HSV2 and syphilis prevalence in female sex workers in Kerman, South-East Iran; using respondent-driven sampling. Iran J Public Health. 2012;41(12):60.
- 69. Wong W, Yim Y, Leung T, Lynn H, Ling D. Prevalence and risk factors of sexually transmitted infections in female sex workers in Hong Kong. Hong Kong Med J. 2012;18suppl3:42-6.
- 70. Dunford L, Carr MJ, Dean J, Nguyen LT, Thi THT, Nguyen BT, et al. A multicentre molecular analysis of hepatitis B and blood-borne virus coinfections in Viet Nam. PloS One. 2012;7(6):e39027.
- 71. Wang JJ, Zhu ZB, Xi Y, Jing W, Wang HB, Lin F, et al. Herpes simplex virus type 2 risks in female sex workers in the China-Vietnam border county of Hekou. Biomed Environ Sci. 2012;25(6):706-10.
- 72. Dunford L, Carr MJ, Dean J, Waters A, Nguyen LT, Thi THT, et al. Hepatitis C virus in Vietnam: high prevalence of infection in dialysis and multi-transfused patients involving diverse and novel virus variants. PloS One. 2012;7(8):e41266.
- 73. Cárcamo CP, Campos PE, García PJ, Hughes JP, Garnett GP, Holmes KK, et al. Prevalences of sexually transmitted infections in young adults and female sex workers in Peru: a national population-based survey. e Lancet Infect Dis. 2012;12(10):765-73.
- 74. Barua P, Mahanta J, Medhi GK, Dale J, Paranjape RS, Thongamba G. Sexual activity as risk factor for hepatitis C virus (HCV) transmission among the female sex workers in Nagaland. Indian J Med Res. 2012;136 (7):30.
- 75. Delgado S, Alvarez B, Goins J, Romero L, Acevedo E, Paz-Bailey G, et al. O1-S08. 04 Central American Surveillance Survey of Sexual Behaviour and Prevalence of HIV/STIs in vulnerable populations: female sex workers, Nicaragua, 2009. Sex Transm Infect. 2011;87(Suppl 1):A42-A.
- 76. Qyra S, Basho M, Bani R, Dervishi M, Ulqinaku D, Bino S, et al. Behavioral risk factors and prevalence of HIV and other STIs among female sex workers in Tirana, Albania. New Microbiol. 2011;34(1):105-8.
- 77. Yang Y, Yao J, Gao M, Su H, Zhang T, He N. Herpes simplex virus type 2 infection among female sex workers in Shanghai, China. AIDS Care. 2011;23(sup1): 37-44.
- 78. Kassak K, Mahfoud Z, Kreidieh K, Shamra S, Afifi R, Ramia S. Hepatitis B virus and hepatitis C virus infections among female sex workers and men who have sex with men in Lebanon: prevalence, risk behaviour and immune status. Sex Health. 2011;8(2):229-33.
- 79. Hemalatha R, Kumar RH, Venkaiah K, Srinivasan K, Brahmam G. Prevalence of & knowledge, attitude & practices towards HIV & sexually transmitted infections (STIs) among female sex workers (FSWs) in Andhra Pradesh. Indian J Med Res. 2011;134(4):470.
- 80. Kassaian N, Ataei B, Yaran M, Babak A, Shoaei P. Hepatitis B and C among women with illegal social

- behavior in Isfahan, Iran: Seroprevalence and associated factors. Hepat Mon. 2011;11(5):368.
- 81. Braunstein SL, Ingabire CM, Geubbels E, Vyankandondera J, Umulisa M-M, Gahiro E, et al. High burden of prevalent and recently acquired HIV among female sex workers and female HIV voluntary testing center clients in Kigali, Rwanda. PloS One. 2011;6(9): e24321.
- 82. Znazen A, Frikha-Gargouri O, Berrajah L, Bellalouna S, Hakim H, Gueddana N, et al. Sexually transmitted infections among female sex workers in Tunisia: high prevalence of Chlamydia trachomatis. Sex Transm Infect. 2010;86(7):500-5.
- 83. Todd CS, Nasir A, Stanekzai MR, Bautista CT, Botros BA, Scott PT, et al. HIV, hepatitis B, and hepatitis C prevalence and associated risk behaviors among female sex workers in three Afghan cities. AIDS (London, England). 2010;24(0 2):S69.
- 84. Zermiani M, Frighetto R, Galvan U, Cruciali M, Serpelloni G, editors. Counselling and screening for sexually transmitted diseases in a cohort of prostitutes in the north-east of italy. Infection; 2010: URBAN & VOGEL NEUMARKTER STRASSE 43, D-81673 MUNICH, GERMANY.
- 85. Shethwala ND, Mulla SA, Kosambiya J, Desai VK. Sexually transmitted infections and reproductive tract infections in female sex workers. Indian J Pathol Microbiol. 2009;52(2):198.
- 86. Wang H, Chen RY, Ding G, Ma Y, Ma J, Jiao JH, et al. Prevalence and predictors of HIV infection among female sex workers in Kaiyuan City, Yunnan Province, China. Int J Infect Dis. 2009;13(2):162-9.
- 87. Cavlek TV, Margan IG, Lepej SZ, Kolaric B, Vince A. Seroprevalence, risk factors, and hepatitis C virus genotypes in groups with high-risk sexual behavior in Croatia. J Med Virol. 2009;81(8):1348-53.
- 88. Shahmanesh M, Cowan F, Wayal S, Copas A, Patel V, Mabey D. The burden and determinants of HIV and sexually transmitted infections in a population-based sample of female sex workers in Goa, India. Sex Transm Infect. 2009;85(1):50-9.
- 89. Bautista CT, Pando MA, Reynaga E, Marone R, Sateren WB, Montano SM, et al. Sexual practices, drug use behaviors, and prevalence of HIV, syphilis, hepatitis B and C, and HTLV-1/2 in immigrant and non-immigrant female sex workers in Argentina. J Immigr Minor Health. 2009;11(2):99-104.
- 90. Uusküla A, Fischer K, Raudne R, Kilgi H, Krylov R, Salminen M, et al. A study on HIV and hepatitis C virus among commercial sex workers in Tallinn. Sex Transm Infect. 2008;84(3):189-91.
- 91. Yun H, Park J, Choi I, Kee M, Choi B, Kim S. Prevalence of human papillomavirus and herpes simplex virus type 2 infection in Korean commercial sex workers. J Microbiol Biotechnol. 2008;18(2):350-4.
- 92. Ngo TD, Laeyendecker O, Li C, Tai H, Cui M, Lai S, et al. Herpes simplex virus type 2 infection among commercial sex workers in Kunming, Yunnan Province, China. Int J STD AIDS. 2008;19(10):694-7.

- 93. Xu JJ, Wang N, Lu L, Pu Y, Zhang GL, Wong M, et al. HIV and STIs in clients and female sex workers in mining regions of Gejiu City, China. Sex Transm Dis. 2008;35(6):558-65.
- 94. Forbi J, Onyemauwa N, Gyar S, Oyeleye A, Entonu P, Agwale S. High prevalence of hepatitis B virus among female sex workers in Nigeria. Rev Inst Med Trop Sao Paulo. 2008;50(4):219-21.
- 95. Linhart Y, Shohat T, Amitai Z, Gefen D, Srugo I, Blumstein G, et al. Sexually transmitted infections among brothel-based sex workers in Tel-Aviv area, Israel: high prevalence of pharyngeal gonorrhoea. Int J STD & AIDS. 2008;19(10):656-9.
- 96. Taylor A, Hutchinson SJ, Gilchrist G, Cameron S, Carr S, Goldberg DJ. Prevalence and determinants of hepatitis C virus infection among female drug injecting sex workers in Glasgow. Harm Reduct J. 2008;5(1):11.
- 97. Gul U, Kilic A, Sakizligil B, Aksaray S, Bilgili S. Magnitude of sexually transmitted infections among female sex workers in Turkey. Eur Acad Dermatol Venereol. 2008;22(9):1123-4.
- 98. Barua P, Laskar N, Medhi G, Apum B, Mahanta J. Sexual Transmission of Hepatitis C Virus Among Female Sex Workers in India. Int J Infect Dis. 2008;12: e416-e7.
- 99. Monsalve-Castillo F, Gómez-Gamboa L, Albillos A, Álvarez-Mon M, Costa-León L, Araujo MS, et al. Hepatitis C virus in populations at risk for infection. Venezuela. Rev Esp Enferm Dig. 2007;99(6):315-9.
- 100. Passos ADC, Figueiredo JFdC, Martinelli AdLC, Villanova MG, Nascimento MPd, Gaspar AMC, et al. Hepatitis B among female sex workers in Ribeirão Preto-São Paulo, Brazil. Rev Bras Epidemiol. 2007;10: 517-24.
- 101. Vandepitte JM, Malele F, Kivuvu D-M, Edidi S, Muwonga J, Lepira F, et al. HIV and other sexually transmitted infections among female sex workers in Kinshasa, Democratic Republic of Congo, in 2002. Sex Transm Dis. 2007;34(4):203-8.
- 102. Davies SC, Taylor JA, Sedyaningsih-Mamahit ER, Gunawan S, Cunningham AL, Mindel A. Prevalence and risk factors for herpes simplex virus type 2 antibodies among low-and high-risk populations in Indonesia. Sex Transm Dis. 2007;34(3):132-8.
- 103. Papadogeorgaki H, Caroni C, Frangouli E, Flemetakis A, Katsambas A, Hadjivassiliou M. Prevalence of sexually transmitted infections in female sex workers in Athens, Greece— 2005. Eur J Dermatol. 2006;16(6):662-5.
- 104. Theng T, Sen P, Tan H, Wong M, Chan K. Seroprevalence of HSV-1 and 2 among sex workers attending a sexually transmitted infection clinic in Singapore. Int J STD & AIDS. 2006;17(6):395-9.
- 105. Pando MA, Berini C, Bibini M, Fernandez M, Reinaga E, Maulen S, et al. Prevalence of HIV and other sexually transmitted infections among female comer-cial sex workers in Argentina. Am J Trop Med Hyg. e2006; 74(2):233-8.
- 106. Nigro L, Larocca L, Celesia BM, Montineri A, Sjoberg J, Caltabiano E, et al. Prevalence of HIV and

- other sexually transmitted diseases among Colombian and Dominican female sex workers living in Catania, Eastern Sicily. J Immigr Minor Health. 2006;8(4):319-23
- 107. Dolar N, Serdaroglu S, Yilmaz G, Ergin S. Seroprevalence of herpes simplex virus type 1 and type 2 in Turkey. J Eur Acad Dermatol Venereol. 2006;20 (10):1232-6.
- 108. Kweon S-S, Shin M-H, Song H-J, Jeon D-Y, Choi J-S. Seroprevalence and risk factors for hepatitis C virus infection among female commercial sex workers in South Korea who are not intravenous drug users. Am J Trop Med Hyg. 2006;74(6):1117-21.
- 109. Fox J, Taylor G, Day S, Parry J, Ward H. How safe is safer sex? High levels of HSV-1 and HSV-2 in female sex workers in London. Epidemiol Infect. 2006;134(5): 1114-9.
- 110. Cwikel JG, Lazer T, Press F, Lazer S. Sexually transmissible infections among illegal female sex workers in Israel. Sex Health. 2006;3(4):301-3.
- 111. O'Farrell N, Vu Thuong N, Van Nghia K, Thu Tram L, Thanh Long N. HSV-2 antibodies in female sex workers in Vietnam. Int J STD & AIDS. 2006;17(11): 755-8.
- 112. Didelot-Rousseau M-N, Nagot N, Costes-Martineau V, Valles X, Ouedraogo A, Konate I, et al. Human papillomavirus genotype distribution and cervical squamous intraepithelial lesions among high-risk women with and without HIV-1 infection in Burkina Faso. Br J of Cancer. 2006;95(3):355.
- 113. Inciardi JA, Surratt HL, Kurtz SP. HIV, HBV, and HCV infections among drug-involved, inner-city, street sex workers in Miami, Florida. AIDS Behav. 2006;10 (2):139-47.
- 114. Chen X-s, Yin Y-P, Liang G-J, Gong X-D, Li H-S, Poumerol G, et al. Sexually transmitted infections among female sex workers in Yunnan, China. AIDS Patient Care & STDS. 2005;19(12):853-60.
- 115. del Amo J, González C, Losana J, Clavo P, Munoz L, Ballesteros J, et al. Influence of age and geographical origin in the prevalence of high risk human papillomavirus in migrant female sex workers in Spain. Sex Transm Infect. 2005;81(1):79-84.
- 116. Nigam P, Shrivastava S. Letter to Editor-HIV and HbsAg seroprevalence in commercial sex workers in Raipur (Chhattisgarh) area. Indian J Dermatol Venerel Leprol. 2005;71(1):52-3.
- 117. Nessa K, Waris S-A, Sultan Z, Monira S, Hossain M, Nahar S, et al. Epidemiology and etiology of sexually transmitted infection among hotel-based sex workers in Dhaka, Bangladesh. J Clin Microbiol. 2004; 42(2):618-21.
- 118.Belza M, Clavo P, Ballesteros J, Menendez B, Castilla J, Sanz S, et al. Social and work conditions, risk behavior and prevalence of sexually transmitted diseases among female immigrant prostitutes in Madrid (Spain). Gac Sanit. 2004;18(3):177-83.
- 119. Taketa K, Ikeda S, Suganuma N, Phornphutkul K, Peerakome S, Sitvacharanum K, et al. Differential seroprevalences of hepatitis C virus, hepatitis B virus

- and human immunodeficiency virus among intravenous drug users, commercial sex workers and patients with sexually transmitted diseases in Chiang Mai, Thailand. Hepatol Res. 2003;27(1):6-12.
- 120. Uribe-Salas F, Conde-Glez C, Juarez-Figueroa L, Hernandez-Castellanos A. Socio-demographic characteristics and sex practices related to herpes simplex virus type 2 infection in Mexican and Central American female sex workers. Epidemiol Infect 2003; 131(2):859-65.
- 121. Miyazaki M, Babazono A, Kato M, Takagi S, Chimura H, Une H. Sexually transmitted diseases in Japanese female commercial sex workers working in massage parlors with cell baths. J Infect Chemother. 2003;9(3):248-53.
- 122. Qutub M, Akhter J. Epidemiology of genital herpes (HSV-2) among brothel based female sex workers in Bangladesh. Eur J Epidemiol 2003;18(9):903-5.
- 123. Valdivia J, Rivera S, Ramírez D, Bussalleu A, Huerta-Mercado J, Pinto J, et al. Hepatitis C virus infection in female sexual workers from northern Lima. Rev Gastroenterol Peru. 2003;23(4):265-8.
- 124. Risbud A, Pereira M, Meherdale S, Gangakhedkar R, Ghate M, Joshi S, et al. Lack of evidence for sexual transmission of hepatitis C virus in patients attending STD clinics in Pune, India. Sex Transm Infect. 2003; 79(5):425.-
- 125. Huo TI, Wu JC, Wu SI, Chang AL, Lin SK, Pan CH, et al. Changing seroepidemiology of hepatitis B, C, and D virus infections in high-risk populations. J Med Virol. 2004;72(1):41-5.
- 126. Camejo MI, Mata G, Díaz M. Prevalence of hepatitis B, hepatitis C and syphilis in female sex workers in Venezuela. Rev Saude Publica. 2003;37(3): 339-44.
- 127. Bystricka M, Krikova Z, Kŕcová M, Koncova K, Kovacova E, Vaculikova A, et al. Sexually transmitted infections among prostitutes in Bratislava, Slovakia. Acta Virol. 2003;47(3):167-72.
- 128. Uribe-Salas F, Conde-Glez CJ, Juarez-Figueroa L, Hernandez-Castellanos A. Sociodemographic dynamics and sexually transmitted infections in female sex workers at the Mexican–Guatemalan border. Sex Transm Dis. 2003;30(3):266-71.
- 129. Miyazaki M, Takagi S, Kato M, Une H. Prevalences of and risk factors for sexually transmitted diseases among Japanese female commercial sex workers in middle-and high-class soaplands in Japan. Int J STD & AIDS. 2002;13(12):833-8.
- 130. Othman BM, Monem FS. Prevalence of heaptitis C virus antibodies among intravenous drug abusers and prostitutes in Damascus, Syria. Saudi Med J. 2002;23 (4):393-5.
- 131. Laurent C, Henzel D, Mulanga-Kabeya C, Maertens G, Larouze B, Delaporte E. Seroepidemiological survey of hepatitis C virus among commercial sex workers and pregnant women in Kinshasa, Democratic Republic of Congo. Int J Epidemiol. 2001; 30(4):872-7.

- 132. Ishi K, Suzuku F, Saito A, Yoshimoto S, Kubota T. Prevalence of human immunodeficiency virus, hepatitis B and hepatitis C virus antibodies and hepatitis B antigen among commercial sex workers in Japan. Infect Dis Obstet Gynecol. 2001;9(4):215-9.
- 133. Rahman M, Alam A, Nessa K, Hossain A, Nahar S, Datta D, et al. Etiology of sexually transmitted infections among street-based female sex workers in Dhaka, Bangladesh. J Clin Microbiol. 2000;38(3):1244-6
- 134. Limpakarnjanarat K, Mastro TD, Saisorn S, Uthaivoravit W, Kaewkungwal J, Korattana S, et al. HIV-1 and other sexually transmitted infections in a cohort of female sex workers in Chiang Rai, Thailand. Sex Transm Infect.1999;75(1):30-5.
- 135. Ghebrekidan H, Rudén U, Cox S, Wahren B, Grandien M. Prevalence of herpes simplex virus types 1 and 2, cytomegalovirus, and varicella-zoster virus infections in Eritrea. J Clin Virol.1999;12(1):53-64.
- 136. Conde-Glez CJ, Juárez-Figueroa L, Uribe-Salas F, Hernández-Nevárez P, Schmid DS, Calderón E, et al. Analysis of herpes simplex virus 1 and 2 infection in women with high risk sexual behaviour in Mexico. Int J Epidemiol.1999;28(3):571-6.
- 137. Suganuma N, Ikeda S, Taketa K, Wang DH, Yamamoto H, Phornphukutkul K, et al. Risk analysis of the exposure to GB virus C/hepatitis G virus among populations of intravenous drug users, commercial sex workers and male outpatients at STD clinic in Chiang Mai, Thailand: a cross-sectional case-control study. Acta Med Okayama.1998;52(3):161-7.
- 138. Juárez-Figueroa L, Uribe-Salas F, Conde-Glez C, Hernández-Avila M, Olamendi-Portugal M, Uribe-Zúñiga P, et al. Low prevalence of hepatitis B markers among Mexican female sex workers. Sex Transm Infect.1998;74(6):448-50.
- 139. Dada AJ, Ajayi A, Diamondstone L, Quinn TC, Blattner WA, Biggar RJ. A serosurvey of Haemophilus ducreyi, syphilis, and herpes simplex virus type 2 and their association with human immunodeficiency virus among female sex workers in Lagos, Nigeria. Sex Transm Infect.1998;25(5):237-42.
- 140. Ghebrekidan H, Cox S, Wahren B, Grandien M. Prevalence of infection with HIV, hepatitis B and C viruses, in four high risk groups in Eritrea. Clin Diag Virol.1998;9(1):29-35.
- 141. McDonnell R, McDonnell P, Neill M, Mulcahy F. Health risk profile of prostitutes in Dublin. Int J STD & AIDS.1998;9(8):485-8.
- 142. Uribe-Salas F, Hernandez-Avila M, Conde-González C, Juárez-Figueroa L, Allen B, Anaya-Ocampo R, et al. Low prevalences of HIV infection and sexually transmitted disease among female commercial sex workers in Mexico City. Am J Public Health. 1997; 87(6):1012-5.
- 143. Tsakris A, Kyriakis K, Chryssou S, Papoutsakis G. Seroprevalence of hepatitis B markers among female and transsexual sex workers in Athens, Greece. Genitourin Med. 1997;73(3):223.

- 144. Luksamijarulkul P, Deangbubpha A. Hepatitis C antibody prevalence and risk factors of some female sex workers in Thailand. Age (years).1997;20(49):24.50.
- 145. Mesquita PE, Hernandez Granato CF, Castelo A. Risk factors associated with hepatitis C virus (HCV) infection among prostitutes and their clients in the city of Santos, Sao Paulo State, Brazil. J Med Virol. 1997;51 (4):338-43.
- 146. Sattar H, Islam M. Hepatitis B virus markers among the prostitutes of Dhaka. Bangladesh Med Res Counc Bull. 1996;22(1):8-11.
- 147. Nakashima K, Kashiwagi S, Hayashi J, Urabe K, Minami K, Maed Y. Prevalence of hepatitis C virus infection among female prostitutes in Fukuoka, Japan. J Gastroenterol.1996;31(5):664-8.
- 148. Weber B, Rabenau H, Berger A, Scheuermann E, Staszewski S, Kreuz W, et al. Seroprevalence of HCV, HAV, HBV, HDV, HCMV and HIV in high risk groups/Frankfurt a. M., Germany. Zentralbl Bakteriol. 1995;282(1):102-12.
- 149. Watts DM, Corwin AL, Omar MA, Hyams KC. Low risk of sexual transmission of hepatitis C virus in Somalia. Trans R Soc Trop Med Hyg. 1994;88(1):55-6.
- 150. Gotuzzo E, Sánchez J, Escamilla J, Carrillo C, Phillips IA, Moreyra L, et al. Human T cell lymphotropic virus type I infection among female sex workers in Peru. J Infect Dis. 1994;169(4):754-9.
- 151. Develoux M, Meynard D, Dupont A, Delaporte E. Hepatitis C virus antibodies in prostitutes in Niger. Trans R Soc Trop Med Hyg 1994;88(5):536.
- 152. Wu JC, Lin HC, Jeng FS, Ma GY, Lee SD, Sheng WY. Prevalence, infectivity, and risk factor analysis of hepatitis C virus infection in prostitutes. J Med Virol. 1993;39(4): 312-7.
- 153. Ndumbe PM, Skalsky J. Hepatitis C virus infection in different populations in Cameroon. Scand J Infect Dis 1993;25(6):689-92.
- 154. Echevarria S, San Miguel G, Pelayo T, Salas C, Casafont F, Crespo F, et al. Risk for hepatitis C virus (HCV) and human immunodeficiency virus (HIV) infections among prostitutes. Genitourin Med. 1993;69 (4):321.
- 155. Tuncel E, Çelebi S, Babacan M. The prevalence of hepatitis B in prostitutes working at the Erzurum brothel. Mikrobiyol Bul.1992;26(1):50-5.
- 156. Orduna A, Bratos M, Gutierrez P, Almaraz A, Eiros J, Martin J, et al. Infection by hepatitis B and C virus in non-intravenous drug using female prostitutes in Spain. Eur J Epidemiol.1992;8(5):656-9.
- 157. Gutierrez P, Orduna A, Bratos M, Eiros J, Gonzalez J, Almaraz A, et al. Prevalence of anti-hepatitis C virus antibodies in positive FTA-ABS non-drug abusing female prostitutes in Spain. Sex Transm Dis.1992;19 (1):39-40.
- 158. Wtr J-S, Lu CF, Lin W-TL-S-Y. Prevalence of Antibodies to Hepatitis C Virus (anti-HCV; in Different Populations in Taiwan. Zhonghua Min Guo Wei Sheng Wu Ji Mian Yi Xue Za Zhi. 19;24(1):55-60.

- 159. Lee SD, Chan CY, Wang YJ, Wu JC, Lai KH, Tsai YT, et al. Seroepidemiology of hepatitis C virus infection in Taiwan. Hepatology1991;13(5):830-3.
- 160. Ramachandran S, Ngeow Y. The prevalence of sexually transmitted diseases among prostitutes in Malaysia. Sex Transm Infect. 1990;66(5):334-6.
- 161. Mak R, Plum J, Van Renterghem L. Human immunodeficiency virus (HIV) infection, sexually trans-mitted diseases and HIV-antibody testing practices in Belgian prostitutes. Sex Transm Infect.1990;66(5): 337-41.
- 162. Hyams KC, Escamilla J, Romero RL, Alvarado EM, Giraldo NB, Papadimos TJ, et al. Hepatitis B infection in a non-drug abusing prostitute population in Mexico. Scand J Infect Dis.1990;22(5):527-31.
- 163. Caballero LR, Caballero CR, Caballero IR, López MS, López FV, Guerrero JR, et al. Prevalence and risk factors of hepatitis B in Spanish prostitutes. Epidemiol Infect.1987;99(3):767-74.
- 164. Jama H, Grillner L, Biberfeld G, Osman S, Isse A, Abdirahman M, et al. Sexually transmitted viral infections in various population groups in Mogadishu, Somalia. Sex Transm Infect.1987;63(5):329-32.
- 165. Goh C, Kamarudin A, Chan S, Rajan V. Hepatitis B virus markers in prostitutes in Singapore. Sex Transm Infect.1985;61(2):127-9.
- 166. De Hoop D, Anker W, Van Strik R, Masurel N, Stolz E. Hepatitis B antigen and antibody in the blood of prostitutes visiting an outpatient venereology department in Rotterdam. Sex Transm Infect.1984;60(5):319-22.
- 167. Kaklamani E, Kyriakidou A, Trichopoulos D, Papoutsakis G, Koumandaki I, Karalis D. Hepatitis B serology in Greek prostitutes: significance of the different serum markers. Epidemiol Infect. 1980;84(2): 257-61.
- 168. Frösner GG, Buchholz HM, GERTH H-J. Prevalence of hepatitis B antibody in prostitutes. Am J Epidemiol 1975;102(3): 241-50.
- 169. Wang H, Men P, Xiao Y, Gao P, Lv M, Yuan Q, et al. Hepatitis B infection in the general population of China: a systematic review and meta-analysis. BMC Infect Dis. 2019;19(1): 811.
- 170. Leroi C, Adam P, Khamduang W, Kawilapat S, Ngo-Giang-Huong N, Ongwandee S, et al. Prevalence of chronic hepatitis B virus infection in Thailand: a systematic review and meta-analysis. Int J Infect Dis. 2016;51:36-43.
- 171. MacLachlan JH, Cowie BC. Hepatitis B virus epidemiology. Cold Spring Harb Perspect Med. 2015;5 (5):a021410.
- 172. Mahmud S, Akbarzadeh V, Abu-Raddad LJ. The epidemiology of hepatitis C virus in Iran: systematic review and meta-analyses. Sci Rep 2018; 8(1): 150.
- 173. Gao Y, Yang J, Sun F, Zhan S, Fang Z, Liu X, et al., editors. Prevalence of anti-HCV antibody among the general population in Mainland China between 1991 and 2015: a systematic review and meta-analysis. Open Forum Infect Dis. 2019;6(3):ofz040.
- 174. Bigna JJ, Amougou MA, Asangbeh SL, Kenne AM, Nansseu JR. Seroprevalence of hepatitis C virus

infection in Cameroon: a systematic review and metaanalysis. BMJ Open 2017;7(8):e015748.

175. Grasso MA, Schwarcz S, Galbraith JS, Musyoki H, Kambona C, Kellogg TA, et al. Estimating the prevalence and predictors of incorrect condom use among sexually active adults in Kenya: results from a nationally representative survey. Sex Transm Dis 2016; 43(2):87.

176. Mellor R, Lovell A. The lived experience of UK street-based sex workers and the health consequences: an exploratory study. Health Promot Int. 2011;27(3): 311-22.

177. Ybarra ML, Mitchell KJ. A national study of lesbian, gay, bisexual (LGB), and non-LGB youth sexual behavior online and in-person. Arch Sex Behav. 2016;45(6):1357-72.

178. Lange WR, Cone EJ, Snyder FR. The association of hepatitis delta virus and hepatitis B virus in parenteral

drug abusers. 1971 to 1972 and 1986 to 1987. Arch Intern Med. 1990;150(2):365-8.

179. He N, Cao H, Yin Y, Gao M, Zhang T, Detels R. Herpes simplex virus-2 infection in male rural migrants in Shanghai, China. Int J STD & AIDS. 2009;20(2): 112-4.

180. Freeman EE, Weiss HA, Glynn JR, Cross PL, Whitworth JA, Hayes RJ. Herpes simplex virus 2 infection increases HIV acquisition in men and women: systematic review and meta-analysis of longitudinal studies. AIDS. 2006;20(1):73-83.

181. Schacker T, Ryncarz AJ, Goddard J, Diem K, Shaughnessy M, Corey L. Frequent recovery of HIV-1 from genital herpes simplex virus lesions in HIV-1–infected men. JAMA. 1998;280(1):61-6