

## SEQUENCE VARIATION IN MEASLES VIRUS F GENE OF AIK-C VACCINE STRAIN AFTER PASSAGE IN VERO CELLS

Zohreh-Azita Sadigh<sup>1,2</sup>, Mahmoud Shamsi Shahrabadi<sup>\*1</sup>, Abbas Shafiyi<sup>2</sup>, Bijan Bamba<sup>3</sup>

*Virology Department of Iran University of Medical Sciences<sup>1</sup>, Razi Vaccine and Serum Research Institute<sup>2</sup>,  
National Institute of Genetic Engineering and Biotechnology<sup>3</sup>.*

---

**Abstract:** The live attenuated AIK-C measles virus which is used for vaccine to control measles in the world has considerably reduced the incidence of this disease. In the present study we have cloned and analyzed the sequence of the cDNA coding for the F protein of AIK-C Vaccine Strain of Measles Virus after limited passage in Vero Cells. Comparison studies demonstrated a high degree of nucleotide sequence homology (98%) to the AIK-C vaccine strain. Altogether four mutations were observed compared to the sequence of AIK-C vaccine strain which was reported by Mori et al and deposited in GeneBank. These mutations caused amino acid changes at position 278, 362, 419 and 549. Change from leucine to phenylalanine at position 278 of the protein that was an early event during adaptation to Vero cells, and also changes at position 362 and 419 were reported by previous study but change at position 549 wasn't report before, and must be study more.

---

**Keywords:** • MeaslesVirus • AIK-C vaccine strain • F gene • pDONR221 • Restriction Endonuclese.

### Introduction

Measles virus is a member of the order Mononegavirales, family Paramyxoviridae, genus Morbillivirus with a non segmented negative strand RNA. The 16 kb linear RNA encoding six structural (N,P,L,H,F,M) and two non structural (C,V) proteins [1]. Two glycoproteins of hemagglutinin (H) and fusion (F) are present as envelope spikes and play a vital role in production of neutralizing antibody. The F is a type I membrane protein that is glycosylated and initially synthesized as the F0 precursor. F0 is transported to Golgi apparatus where it is cleaved by a host protease into two disulfide-linked subunits, F1 and F2. This protein is responsible for virus penetration through fusion of viral and host cell membrane, cell fusion and hemolysis. Induction of antibody responses to F protein is essential for protection [1,2].

Today, the live attenuated vaccines is used to control measles in the world and has dramatically

reduced the incidence of this disease. Many developing countries still have intense MV transmission, especially in children less than 1 year of age. In these countries measles claims the lives of an estimated 700,000 infants and young children annually [3]. The measles vaccine strain AIK-C is one of the safest vaccine virus which was developed from the Edmonston strain through plaque cloning in sheep kidney cells and chick embryonic cells at 33°C [4] and is widely used for prevention of measles infection. Due to high effectiveness and high sero-conversion in young infants, vaccine manufacturers have attempted to produce the AIK-C measles vaccine in large scale using Vero cells. The sequential passage in these cells may have caused some changes in genome nucleotides of the virus [5].

pDONR vectors that were used in our experiment were adapted vectors designed to generate attL-flanked entry clones containing a gene of interest following recombination with an attB PCR product. This vector was used for generating the expression clone by performing an LR reaction between the entry clone and a destination vector of choice like the baculovirus.

---

\* **Corresponding Author:** Mahmoud Shamsi Shahrabadi, Virology Department of Medical Sciences, Rassule Akram Hospital, Nyayesh Ave, Sattarkhan St, Post Code 144561313, Tehran, Iran

The pDONR vectors contain the following elements *rrnB* T1 and T2 transcription terminators, M13 Forward and M13 Reverse priming sites for sequencing of the interest, the recombination sites (attP1 and attP2) derived from DNA recombination sites of bacteriophage  $\lambda$  that allows recombinational cloning of the gene of interest from attB PCR product, *ccdB* and Chloramphenicol resistance genes located between two attP sites for negative and counter selection respectively, Kanamycin resistance gene for selection in *E. coli* and pUC origin for replication and maintenance of the plasmid in *E. coli* [6]. In the present study we have cloned and analyzed the sequence of the DNA coding for F protein of AIK-C strain of MV after three passages in Vero cells.

## Material and methods

### Cells and virus

Vero cells were grown in Dulbecco's modified Eagle Medium supplemented with 5% Fetal Calf Serum and antibiotics and propagated at 37°C. Monolayer of cells were inoculated with Measles Virus AIK-C vaccine strain provided by Dr. Abbas Shafyi (Razi Vaccine and Serum Research Institute), and inoculated at 37°C temperature. When cytopathic effect was observed, infected cells were harvested and stored at -70°C.

### RNA Extraction

Infected cell cultures were centrifuged at low speed for 20 minutes. The clarified supernatant was ultracentrifuged at 20000 r.p.m for 1 hour. The pellet consisted of concentrated virus was used for RNA extraction. RNA was extracted using RNA extraction kit (RNA FAST manufactured in National Institute of Genetic Engineering and Biotechnology). The extracted RNA was stored at -70°C.

### Primers

Primers used for RT-PCR and PCR were designed based on published MV sequences (GeneBank accession No. S58435) and contain recombinant sites (attB1 and attB2 for Forward and Reverse primers respectively shown with Bolded nucleotides) for cloning into pDONR vector. The sequence of the primers were as follow:

**F-F1: 5'GGGG ACA AGT TTG TAC AAA  
AAA GCA GGC TCA ATG TCC ATG ATG  
GGT CTC 3'**

**F-R11: 5'GGGGAC CAC TTT GTA CAA GAA  
AGC TGG GTA GAG CGA CCT TAC ATA  
GGA TT 3'**

We designed also a forward primer complementary the sequence from 6022 to 6041 for thorough sequencing of the F gene as follow:

**FM<sub>1</sub>: 5' TCC ATG GAC CAA CTA TCT TG 3'**  
Amplification of the F gene:

Extracted RNA was used as template for reverse transcription reaction. The RT-PCR reaction was performed in a total 20  $\mu$ l final volume, using 2.5mM of each dNTPs, 100  $\mu$ mol of each primer, reaction buffer (containing 250mM Tris HCl, 250mM KCl, 20mM MgCl<sub>2</sub>, 50mM DTT), 40U Rnase Inhibitor, 200 U M-Mulv Reverse Transcriptase and nuclease free water. Thermocycler was programmed for one cycle at 70°C for 5 minutes, followed by one cycle at 37°C for 5 minutes and one cycle at 42°C for 60 minutes. The reaction was stopped at 70°C for 10 minutes and then the microtube was chilled on ice [7,8].

PCR test was performed in a total of 50  $\mu$ l final volume, using 1 ng cDNA, 10  $\mu$ mol of each primer, 2.5mM of each dNTPs, 2.5mM MgSO<sub>4</sub> and 2.5 units of *pfu* DNA polymerase [9,10]. Thermocycler was programmed for one cycle at 94°C for 5 minutes followed by thirty cycles at 95°C for 1 minutes, 52°C for 1 minutes, 72°C for 1.2 minutes and one cycle at 72°C for 10 minutes, as a final extension.

PCR product was electrophoresed on 1% (w/v) agarose gel and stained by ethidium bromide and visualized by UV transilluminator.

Plasmid and Construction of F gene Cloning vector:

The pDONR221 plasmid and Omnimax chemically competent *E. coli* (Invitrogen) were used for cloning and sequencing.

The full coding region sequence of F protein (AIK-C Stain) was inserted into pDONR221, using the BP recombination reaction by BP clonase (Invitrogen) [11].

### Transformation

One vial of Omnimax chemically competent *E. coli* was thawed on ice, then 1  $\mu$ l of BP recombination reaction were added into the vial and incubated on ice for 30 minutes. The cells were inoculated for 90 seconds at 42°C and the vial removed and kept in ice for 3 minutes. After that 250  $\mu$ l of S.O.C medium kept at room temperature was added to the vial. The vial was


shaken horizontally at 37°C for 1 hour, then 50µl of the transformation mix was spread on a prewarmed selective plate (LB+Kanamycin) and incubated overnight at 37°C [11].

Screening the Colonies:


Clonies carrying plasmid were picked up and suspended in 5 ml of LB broth with appropriate kanamycin concentration. Plasmids were purified from the bacteria using plasmid purification kit (Roche), and then were analyzed by Restriction Endonuclease digestion (BstX I) and sequenced with M13 primers in both direction by a dye terminator method, using the ABI 3130XL sequencer. The Nebcutter software was used for analyzing the restriction sites in PCR product and plasmids. The sequencing results were compared with other sequences deposited in the GeneBank using the BLAST software[ 12].

Results:

PCR amplification of cDNA from Measles Virus F gene with specific primers generated a fragment with approximately 1700 bp in size (Fig.1). This DNA fragment product of F gene was analyzed by EcoR V digestion. This gene has one restriction site for EcoRV, thus after digestion we had two fragments with sizes of approximately 1000bp and 700bp (Fig 2 Lane1).




**Fig-1:** Amplification of F gene (Measles virus AIK-C strain). Lane1: Amplification of F gene with specific primers (1711bp) Lane2: Negative control. LaneM: DNA Molecular weight Marker(0.07-12.2Kbp).




**Fig-2:** Digestion analysis of F gene with EcoRV. Lane1: Digestion analysis of PCR product of F gene(711bp and 1000bp). Lane M: DNA Molecular weight Marker (0.07-12.2 Kbp)

PCR product of the F gene was cloned into the pDONR221 by Bp recombination reaction, and these constructs were used to transform Omnimax E. coli. The presence of recombinant plasmids in the transformed colonies were screened by selection on LB agar medium containing 50µg/ml Kanamycin. After plasmid purification they were analyzed by BstX I restriction enzyme. This enzyme has 2 restriction sites in the F gene at nucleotides 33 and 1359, but has no restriction site in non-recombinant pDONR221 (Fig.3). Therefore, after digestion we had two fragments with approximately 1300bp and 3000bp in size (Fig.4. Lane 2, 4).




**Fig-3:** Schematic presentation of pDONR 221+ F gene of Measles Virus construct with some restriction sites.



**Fig-4:** Confirmation of cloning of F gene into pDONR via restriction enzyme analysis: Lane1: Undigested pDONR + F plasmid 1 . Lane2: digestion of pDONR221 F plasmid 1 with BstX I(1326bp and 3000bp). Lane 3: Undigested pDONR + F plasmid 2. Lane 4: digestion of pDONR221 + F plasmid 2 with BstX I(1326bp and 3000bp). Lane M: DNA Molecular weight marker(0.07-12.2Kbp).

The recombinant plasmids were analyzed by sequencing and the results were compared with the sequences of AIK-C strain of attenuated Measles Virus in the GeneBank 9S58435, Mori et al 1993)[13] using the BLAST Software [12]. Nucleotide comparison showed 4 differences between original AIK-C vaccine strain and AIK-C vaccine strain which passaged in Vero cells (Fig.5).



**Fig-5:** Sequence comparison of measles virus AIK-C vaccine strain (subject) and AIK-C vaccine strain which were passaged 3 times in Vero cells. (Query).

These nucleotide substitution led to 4 amino acid substitution in the F protein (Leucine to phenylalanine at position 278; F278-Phe, Serine to Tyrosine at position 362; F362-Tyr, Histidine to Asparagine at position 419; F419-Asn and serine to Leucine at position 549; F549-Leu).

### Discussion

Most RNA viruses are considered to be a mixture of heterogeneous viruses because of their high incidence of mutation in the RNA genome in nature [14]. Certain viruses such as measles are antigenically stable, whereas others readily give rise to antigenic variants [15]. Analyses of the MV genes of a large number of strains have established that the F gene is highly stable, and up to 1989 only three amino acid changes have been observed in the Edmonston strain and related isolates [16]. The measles vaccine strain AIK-C which was developed from the Edmonston strain which has been attenuated in lamb kidney cells and cloned in chick fibroblast cells at 33°C [4], are widely used for prevention of measles infection in the world [5]. Because of the high effectiveness of vaccination program and low incidence of adverse reaction and high sero-conversion in young infants, a vaccine manufacturer decided to produce the AIK-C measles vaccine in large scale using Vero cells [15]. In this regard a group of investigators studied the genetic stability of the virus and reported that there was three amino acid changes in the F gene [17]. In this study AIK-C vaccine strain was passaged 3 times in Vero cells in order to amplify the F gene. We observed 4 amino acid substitution which were different from those reported by previous studies [17]. A likely explanation of these differences lies in the types of tools and methods used to analyze the virus genome. Another difference lay in the passage history of the sequenced AIK-C viruses, because here the virus was passaged three times in Vero cells, whereas Mori et al used chicken embryo cells [18]. Change from leucine to phenylalanine at position 278 of the F protein was the first mutation occurred during the viral passage in Vero cells. The extent of these genome variation on viral properties remains to be determined.

## References

- 1 Griffin, D.E, 2001. Measles virus, p.1401-1428. In B.N. Fields, O.M Knipe , P.M. Howely, Fields virology 4<sup>th</sup> ed Lippincott Williams and Wilkins publishers.
- 2 Atabani, S.F.,Obeid,O.E.,Chargelegue,D.,Abbay,P.,Whittle,H., Stward, M.W.1997. Identification of an Immunodominant neutralizing and protective epitope from measles virus fusion protein by using human sera from acute infection J.virol 71:7240-7245.
- 3 Premenko – Lanier, Hodag, G.,Rota, P., Tamin, A.,Bellini,W.,McChesney,M.2006. Maternal antibody inhibits both cellular and humoral immunity in response to measles vaccination at birth. Virology.2006 Jul 5;350(2):429-32.
- 4 Sasaki. K.1974. studies on the modification of the live AIK measles vaccine I- adaptation of the further attenuated AIK measles virus (The strain of AIK-L33) to chick embryo cells. Kitasato Arch. Exp.Med .47:1-12.
- 5 Komase,K., Nakayama,T., Iijima,M., Miki,K., Kawanishi,R., Uejima,H. 2006. The phosphoprotein of attenuated measles AIK-C vaccine strain contributes to its temperature-sensitive phenotype.Vaccine.24:826-834.
- 6 Gateway pDONR vectors.Version C .15 Jul 2004.
- 7 Protocol for First Strand cDNA Synthesis( PCR template generation with M-MuLV Reverse Trascriptase) Fermentas.
- 8 Verma,I.M.1981.Reverse transcriptase.The Enzymes (Boyer,P.D.,ed), Academic Press Inc., Vol.14:87-103.
- 9 Protocol for PCR with *pfu* DNA polymerase Fermentas.
- 10 Lundberg, K.S.,et al.1991. High- fidelity amplification using a thermostable DNA polymerase isolated from *Pyrococcus furiosus*,Gene,108:1-6.
- 11 Gateway Technology with Clonase II.Version A. 24 June 2004 Invitrogen.
- 12 Atschul,S.F., Madden,T.L.,Schaffer, A.A., Zhang,J.,Zhang,Z.,Miller.W.,Lipman,D. 1997."Gapped BLAST and PSI-BLAST:a new generation of protein database search programs",Nucleic Acids Res.25:3389-3402.
- 13 Mori,T., Sasaki,K.,Hohimoto,H.,Makino,S. 1993. Molecular cloning and complete nucleotide sequence of the genomic RNA of the AIK-C strain of attenuated measles virus.Virus Genes.7:67-81.
- 14 Domingo,E., Ruiz-Jarab,CM., Sierra,S.,Arias,a., Pariente,N., Baranowski,E. 2002. Mini Review. Emergence and selection of RNA variants: memory and extinction. Virus Res.82:39-44.
- 15 Rota,J.S., Redd,S.B.,Redd,s.c.,Pattamadilok,S.,Bellini,W.J.1998. Genetic analysis of measles viruses isolated in the United States,1995-1996.J.Infect. Dis. 177:204-208.
- 16 Rota,J.S., Hummel,K.B., Rota,P.A., Bellini,W.J. 1992. Genetic variability of the glycoprotein genes of current wild-type measles isolates.Virology.188:135-142.
- 17 Uejima,H., Nakayama,T., Komase,K. 2006. Passage in Vero cells alters the characteristics of measles AIK-C vaccine strain.Vaccine. 24:931-936.
- 18 Parks,C.L.,Lerch,R.A.,Walpita,P.,Wang,H-P.,Sidhu,M.S.,Udem,S.A.2001. Comparison of predicted amino acid sequences of Measles Virus Strains in the Edmonston Vaccine Lneage.J.Virol.75:910-920.