

Original Article

Phylogenetic Analysis of Hemagglutinin Gene of H9N2 Avian Influenza Viruses Isolated from Chicken in Iran in 2010-2011: Emerging of a New Subgroup

Norouzian H^{1*}, Gholami SJ², Vasfi-Marandi M²

1. Department of Clinical Sciences, Faculty of Veterinary Medicine, Lorestan University, Khorram Abad, Iran.
2. Department of Clinical Sciences, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran.

Abstract

Background and Aims: Hemagglutinin (HA) protein of Avian Influenza (AI) plays an essential role in the virus pathogenicity. AI H9N2 subtype causes significant economic loss in broiler and layer in poultry farms in Iran. AI viruses have a great involvement in evolutionary changes at nucleotide and amino acid levels and vaccines could induce faster rates of such changes. Up-dated understanding of the genetic changes of AI viruses circulating in Iran is necessary for controlling AI.

Materials and Methods: Sequence analysis and phylogenetic study of the HA gene of three H9N2 subtype of AI isolates in Iran in 2010-2011 were studied.

Results: Cleavage site of the Iranian 2010-2011 isolates possessed a different motif. Amino acid residue at position 226 at receptor binding site in these isolates was Leucine, which was similar to human viruses. The epitopes for HA showed a great variation related to the year of isolation. According to phylogenetic analysis, Iranian isolates were divided into two main subgroups. But, viruses isolated in this study formed a third minor subgroup. Degree of homology between the 2010-2011 isolates and former Iranian isolates was significantly low.

Conclusion: The results revealed that HA of new Iranian AI H9N2 isolates have undergone extensive genetic changes. Definitely, continuous monitoring of genetic changes is a useful tool for updating control strategy for AI outbreak in Iran.

Keywords: Avian Influenza; H9N2; Hemagglutinin; Phylogenetic analysis; Iran

Introduction

Avian influenza (AI) virus is a member of the family orthomyxoviridae. These viruses contain eight single-stranded RNA segments, which encode ten proteins, including two surface glycoproteins, hemagglutinin (HA) and neuraminidase (NA) (1). Up to now, 16 HA and 9 NA subtypes

have been identified (2). HA protein plays an essential role in virus binding to sialic acid receptor on the host cell surface and fusion of virus to endosomal membrane, which facilitates entry of virus into the host cell (1).

AI H9N2 subtype typically causes mild clinical signs in birds. However, significant mortality in broilers and significant decrease in egg production have been reported due to H9N2 subtype, particularly in co-infection with other respiratory agents (3, 4). H9N2 influenza viruses have become widespread in the poultry farms of the world (4-6). According to the phylogenetic and antigenic analysis, Asian H9N2 viruses are classified as three major

*Corresponding author: Hassan Norouzian. Faculty of Veterinary medicine, Lorestan University, Khorram Abad, Iran, PO Box: 68137-17133. Tel: (+98) 661 62 00 109.

Email: noroozianh@gmail.com

sublineages represented by A/Quail/Hong Kong/G1/97 (G1 group), A/Chicken/Hong Kong/G9/97 (G9 group), and A/chicken/Korea/38349-p96323/96 (Korean group) (7).

To control AI infections in poultry, inactivated vaccines have been widely used in Iran, Pakistan and China (8-10). However, it is revealed that vaccines could induce faster rates of antigenic drift in influenza viruses due to providing a positive selective pressure (9). On the other hands, co-circulation of AI H9N2 viruses with other subtypes, including H5 and H7, may lead to the genetic reassortment in the viruses (11). For these reasons, up-dated understanding of the genetic changes of AI viruses circulating in Iran is important for controlling AI. In this study, the HA gene of three AI H9N2 strains isolated from poultry chicken farms in Iran during 2010 and 2011, as well as, other reported Iranian H9N2 isolates were genetically analyzed and nucleotide changes were evaluated.

Methods

Viruses

Three strains, including :

A/Chicken/Iran/N101/2011(H9N2), A/Chicken/Iran/N102/2011(H9N2) and A/Chicken/Iran/EBGV-88/2010) were isolated from three broiler farms affected by respiratory disease complex (RDC). Initial isolation and subtype identification of the virus was determined by standard methods (12) and PCR tests. All the other H9N2 viral sequences of hemagglutinin genes were present in the GenBank database.

Cell transfection

COS-7 and BHK-21 cell lines were received from the National Cell Bank of Iran (NCBI) affiliated with the Pasteur Institute of Iran, cultured and maintained in Dulbecco's modified Eagle's Medium (DMEM, Gibco) supplemented with 10% fetal bovine serum (FBS, Gibco). Briefly, one day before transfection, cells from confluent culture were transferred to 24 well plate (Nunc, Roskilde, Denmark) containing 10^5 cells per well. The cells were incubated for 24 h at 37°C in a

humidified incubator containing 5% CO₂ to achieve approximately 70-75% confluency on the day of transfection. Each well of cells was transfected with 0.8µg of plasmid DNA encoding the HA1 gene using lipofectamin 2000 reagent (Invitrogen, Carlsbad, CA; Cat. No. 11668-027) according to the manufacture's instruction. After 48h incubation at 37°C, cells were examined by fluorescence microscope.

RNA Extraction and RT-PCR

Viral RNA was extracted from allantoic fluid by RNX plus kit (Cinnagen, Iran), according to the manufacturer instruction. Reverse transcription was done by using oligonucleotide influenza universal primer, Uni12. Amplification of the hemagglutinin full-length gene was carried out by PCR as described previously (13) using two pairs of specific primers as follows:

P-HA1-F: 5'-AGC AAA AGC AGG GG-3' and

P-HA1-R: 5'-GTG YCC ATA CCA TGG RGC-3' (1 to 832),

P-HA2-F: 5'- ACC AGG TCA GAC ATT GCG AG-3' and

P-HA2-R: 5'- AGT AGA AAC AAG GGT GTT TTT GC-3' (766-1742).

Gene Sequence

PCR products were subjected to electrophoresis in a 1% (w/v) agarose gel and DNA fragments of the expected length were extracted and purified with GeneJET™ Gel Extraction kit (Fermentas, Canada, catalogue#k0691). The purified DNA fragments were cloned into pTZ57R/T cloning vector (Fermentas, Canada, catalog#k1213). Three clones of each fragment were sequenced using M13 forward and M13 reverse (promega, USA) at Sequetech Co. Ltd., USA. The sequences identified of at least two clones were kept for further usage.

Sequence Analysis and Phylogenetic Study

Pairwise sequence alignments were performed using the Clustal W alignment algorithm. Sequence similarity and phylogenetic relationship of different H9N2 subtype viruses was performed with MegAlign program (DNASTAR Inc., Madison, WI, USA).

Nucleotide Sequence Accession Numbers

Table 1. Amino acid substitutions in the cleavage site, Receptor binding site and PGS of H9 HAs of Iranian H9N2 isolates.

virus	Accession no.	Cleavage site	Receptor binding site			Potential Glycosilation Sites						
			190	226	227	105	166	206	218	283	492	551
11T/99	AF508558	RSSR	A	Q	Q	+	-	-	-	-	+	+
B308A/04	EF063728	RSSR	A	L	Q	+	-	-	-	-	+	+
EBGV-88/10	JN646748	KSSR	T	L	I	+	-	-	-	-	+	+
N101/11	JQ970436	KSSR	T	L	I	+	-	-	-	-	+	+
N102/11	JQ970437	KSSR	A	L	I	+	-	-	-	-	+	+
G1/97	AF156378	RSSR	E	L	Q	+	-	+	+	-	+	

The sequences determined in this study are available in the GenBank under accession numbers: JQ970436, JQ970437 and JN646748.

Results

Analyses of the hemagglutinin sequence of the Iranian H9N2 isolates

In this study, 1742 nucleotides of the HA gene of the three isolates (A/Chicken/Iran/N101/2011(H9N2), Chicken/Iran/N102/2011(H9N2) and A/Chicken/Iran/EBGV-88/2010) were sequenced and amino acid sequences of the HA were deduced from the nucleotide sequence.

HA cleavage site motif sequence of the most Iranian isolates was: -R-S-S-R*G-. Amino acids of cleavage site of the isolates sequenced in this study possessed a -K-S-S-R*G- motif. Some other substitutions have happened in

some Iranian isolates. Four isolates harbor -R-S-N-R*G- at the same position (Table 1).

In the HA protein, the amino acids at positions 183, 190, 226, 227 and 228 (H3 numbering) are major receptor binding sites (RBS). Amino acid residue at position 226 in isolates sequenced in this study, was Leucine. In Iranian H9N2 viruses, the number of those possessed the human H9N2 virus-like motif, 226Leu, has increased after 2004. There is a substitution at position 227, Glu→Ile, in almost all Iranian H9N2 viruses isolated after 2004, including three viruses sequenced in this study. In one of the viruses sequenced in this study (A/Chicken/Iran/N101/2011), amino acid residue Thr replaced Ala at position 190. In Iranian isolates, substitutions in this position have happened, which Ala replaced by Thr, Ile and Val in some recent isolates (Table 1).

Potential glycosilation site (PGS) motifs -N-X-T/S- (X may be any amino acid except proline)

Table 2. Amino acid substitutions in the antigenic sites of H9 HAs of Iranian H9N2 isolates.

virus	Accession no.	Antigenic Sites					
		108	134	154	312	324	431
11T/99	AF508558	+	-	-	+
B308A/04	EF063728	+	-	-	+
EBGV-88/10	JN646748	+	+	.T....	+	+
N101/11	JQ970436	+	+	.I....IG	+	+
N102/11	JQ970437	+	+	.I....IG	+	+

Table 3. Homology of the nucleotide sequences of HA of Iranian isolates belonging to different sublineages.

sublineage	Similarity between viruses (%)								
	First sublineage			Second sublineage			Third sublineage		
	Max	Min	Average	Max	Min	Average	Max	Min	Average
First	100	91.8	96.4	-	-	-	91.7	89.4	90.5
Second	-	-	-	99.9	93.7	97.4	93.8	91.1	92.8
Third ¹	-	-	-	-	-	-	98.7	99.1	98.9

in the HA molecule of viruses sequenced in this study and most other Iranian H9N2 isolates were seven sites at positions 29, 105, 141, 298 and 305 in HA1 and 492 and 551 in HA2 region, C-terminal to the membrane anchor. However, some of these PGSs have been lost in some isolates. On the other hand, some Iranian H9N2 isolates possessed new PGSs at positions 158, 198, 210 and 275 due to point mutations (Table 1).

The epitopes for HA showed a great variation related to the year of isolation. The epitope (SFYRNMRWL) from positions 154-162 and the epitope (KAIGLRNV) from positions 324-331 that were mostly absent in viruses isolated between 1998 and 2004, appeared in H9N2 viruses after 2004. Another epitope with high variation at position 312-320, which in most of the viruses isolated before 2004 was (GNCPKYVRV) and after that year it has changed to (GTCPKYIGV). This epitope in two of isolates sequenced in this study was changed to (GICPKYIGV). Some of the epitopes (position 431-439 (AYNAELLVL) and position 134-142 (IFPDTIWNV)) are present in all Iranian H9N2 isolates. In the epitope (CYPGNVENL), positions 108-116, there is an amino acid substitution at position 113 (Val113Ile) in a few Iranian isolates (table 2).

Phylogenetic analyses

The phylogenetic relationship between Iranian isolates and H9N2 viruses isolated in various part of Middle East countries were established. The analysis was based on the nucleotide sequence of full segment of HA. All of the Iranian isolates fell into the same main group (G1 sublineage), as the viruses isolated in Pakistan, Saudi Arabia, Japan and Israel (Fig 1a). The Iranian isolates can be further divided

into two main subgroups. The first subgroup (11-T like sublineage) included viruses that were isolated between 1998 and 2004. The second subgroup comprised the strains isolated after 2004. Though, some exceptions were seen in both subgroups. Interestingly, three viruses isolated in this study formed a third minor subgroup (Fig 1b). Other Middle East isolates also clustered in two subgroups, with some exceptions, according to the country and time of isolation. Three viruses isolated in this study, placed near the new Pakistan isolates in a different subgroup (Fig 1a).

Analysis of the Iranian isolate nucleotide sequences showed that the difference within the first subgroup was relatively higher than the second subgroup: viruses belonging to the first subgroup exhibited 91.8-100% (average 96.4%) identity and viruses belonging to the second subgroup exhibited 93.7-99.9% (average 97.4%) identity. Three viruses of the third minor subgroup, the isolates sequenced in this study, exhibited higher identity 98.7-99.1% (average 98.9%) (Table3). Interestingly, these isolates also showed relatively high identity to some of the new Pakistan isolates (95%). Degree of homology between these new isolates and former Iranian isolates is relatively low (90.5% and 92.8% for first and second subgroup, respectively). The percent of homology of isolates sequenced in this study with prototype isolate of G1-like sublineage and human H9N2 isolates was 90.9% (Table 3).

Genetic variability amongst the Iranian isolates

In this study, we analyzed 1682 nucleotides encoding 560 amino acids (1-560). Of these 560 amino acids, 537 were characterized as highly conservative amongst the Iranian




Fig. 1. Phylogenetic relationships of HA gene of influenza A viruses isolated in Middle East (a) and Iran (b). The lengths of horizontal lines are proportional to the minimum number of nucleotide differences required to join nodes. Vertical lines are for spacing branches and labels. ck, chicken; qa, Quail; pk, pakistan; Ir, Iran; Is, Israel; Lb, Lebanon; Ind, India; HK, Hong Kong; SA, Saudi Arabia

isolates, whereas the other 23 amino acids showed diverse levels of variability. The subgroup specificity of the substitutions was observed in the following positions: amino acids A¹⁵, S¹⁰¹, A¹²⁶, A¹⁵⁰, S¹⁵⁸, S¹⁶⁶, S¹⁶⁸, S¹⁸³, D²¹⁶, N²¹⁸, M²²⁴, Q²³⁵, G²⁷¹, N³¹³, V³¹⁸, R³¹⁹, and V³²⁷ were typical of the viruses of the first subgroup, whereas T¹⁵, P¹⁰¹, S¹²⁶, S¹⁵⁰, N¹⁵⁸, N¹⁶⁶, L¹⁶⁸, D¹⁸³, N²¹⁶, D²¹⁸, L²²⁴, I²³⁵, E²⁷¹, T³¹³, I³¹⁸, G³¹⁹, and I³²⁷ were typical of those of the second subgroup.

Discussion

Changes in the HA amino acid content at the proteolytic cleavage site (PCS) by either site mutation or insertions of amino acids or both, resulting in an increase in the number of basic amino acids (Arginine and Lysine) at the site provides a template for cleavage by widely distributed furin-like or subtilisin-like endoproteases (11). This is central key to the switch from low to high pathogenicity in H5

and H7 subtypes (14, 15). Nucleotide sequence analysis corresponding to the cleavage site of Iranian isolates revealed no poly-basic motif. Most of the earliest Iranian isolates (since 1998) had the cleavage site motif -RSSR*G- with two basic amino acids, which is rather typical for H9 viruses, including those from Hong Kong (Qa/HK/G1/97 and Dk/HK/Y280/97), Pakistan, Israel, Saudi Arabia and Dubai (16). Some viruses with point mutations of amino acid in the cleavage site (RSNR, RSNK) were isolated on 2007. Then, on 2010 and 2011, all isolated viruses, including viruses sequenced in this study, showed another amino acid substitution (KSSR). This substitution of lysine instead of Arginine at the -4 position of the cleavage site is infrequently observed in H9N2 viruses, but have become fixed in all 2010 and 2011 Iranian and some recent Pakistani isolates (table 1)_(17, 18, 19). The significance, if any, of these mutations is not known. Banks et al. (2000) (16) described 15 amino acid motifs of

the cleavage sites of the H9 influenza viruses, but none of them had more than two basic amino acids. It should be emphasized, that all H9N2 viruses with two or three basic amino acids in the cleavage site have had low pathogenicity for chickens (19, 18). Some researchers suggested that -RSSR*G- motif is the signature of H9N2 LPAI viruses adapted to the chicken host (16, 18, 20). It is interesting that all of these mutations in Iranian isolates have happened after 2007. So, there may be a relationship between time and happening of mutations in AIVs, as observed in Israeli isolates (19).

AI H9N2 viruses have some capabilities to transmit from bird to human and are a candidate for next human flu pandemic (20). The RBS motif of HA is critical for cellular receptor specificity and determining virus host range (21, 22). Glutamine at position 226 indicates a preferential binding to sialic acid (SA) α 2,6-linked to galactose found in avian host and is a major host range determinant for the H2 and the H3 subtypes associated with pandemic human flu infections (23, 24). The substitution of Glu226Leu makes a preferential receptor binding specificity for SA α 2,6-linked galactose, allows H9N2 viruses to replicate about 100-fold more efficiently in human cell culture (23, 25). There is a general increasing trend in the proportion of Iranian H9N2 viruses that possessed the human virus-like motif 226Leu, particularly in viruses isolated after 2004 (Table 1) and in recent Pakistan, UAE and Chinese isolates (7, 18, 26). This result shows increasing potential of recent Iranian H9N2 isolates to transmit to human. The residue at position 190 within the RBS influences the affinity of virus binding to the SA receptor. Amino acids Valine, Threonine and Alanine at this position had high, intermediate and low affinity to SA α 2, 6-linked galactose receptor, respectively (27). Alanine at this position is more typical of viruses isolated from chickens (28). In Iranian isolates, substitutions in this position (A \rightarrow T, A \rightarrow I, A \rightarrow V) have happened in some recent isolates. In one of the viruses sequenced in this study (N101), amino acid Threonine has replaced Alanine, indicating high affinity for

human airway epithelial receptors (Table 1). Nevertheless, the definitive significance of these substitutions is not completely understood and requires detailed structural analysis.

Seven PGSs were present in most of the Iranian H9N2 isolates, including isolates sequenced in this study. However, there were additional or lost PGSs in some of the isolates. Two PGSs at positions 218 and 206, present in representative reference strain of G1-lineage (A/Quail/HK/G1/97) have lost in most of the Iranian isolates (Table 1). The loss or creation of PGSs may represent a selected adaptation of H9N2 viruses transmitted from original host (e.g. Quail) to poultry (14, 29). Besides, these alterations may affect the receptor binding capacity of HA and contribute to the antigenic variations (30, 31). Anyway, the biological significance of these changes in PGSs pattern in the Iranian H9N2 isolates remains to be studied.

A study on evolution of AI H9N2 viruses in the Middle East and Indian sub-continent showed variability in antigenic sites of HA during 1994-2009 (20). Antigenic sites for HA showed a great deal of variation. Observation revealed that epitopes (SFYRNMRWL) from positions 154-162 and (KLAIGLRNV) from positions 324-332 that was absent before 2004 in first subgroup of Iranian isolates, appeared in the second subgroup after 2004 (Table 2). This finding is in agreement with phylogenetic analysis results. In the other epitopes, also, there are some variations between Iranian H9N2 isolates (Table 2). These results suggest that NA epitopes could have evolved due to point mutations and reassortment. Such changes would affect virus antigenicity and pathogenicity of the virus. It is revealed that vaccines could induce faster rates of antigenic drift in human and avian influenza viruses possibly due to inducing a positive selective pressure in the host (32, 33). Besides, there is a concern about that the antigenic differences between the vaccine strain and prevailing H9 viruses might worsen the continued antigenic drift of the viruses. Most of the isolated H9N2 viruses in china after 2003 were antigenically heterologous with the vaccine isolate due to

antigenic drift and commercial inactivated vaccine could not protect chickens against some of the H9N2 viruses in cases of virus shedding or appearing of clinical signs (34). It is better to compare the antigenic properties of HA protein of new H9N2 isolates with vaccine strain and re-evaluate of vaccine efficacy in Iran.

Previous studies on phylogenetic analysis of HA gene have determined that all Iranian H9N2 viruses fell into a special group and came from a single progenitor (13, 20, 35, 36). On the other hands, all Iranian isolates fell into the same main group (G1 sublineage) as the viruses isolated in Pakistan, Saudi Arabia, UAE, India and Israel, indicating they might have originated from a common source (Fig 1a). However, Middle East isolates showed a relationship that may indicate some ecological barriers to virus movement based around geographic location, with isolates grouped according to country of isolation (20). The Iranian isolates can be further divided into two main subgroups (Fig 1b). The first subgroup (11-T like subgroup) viruses isolated mainly between 1998 and 2004 and the second subgroup mainly after 2004, with some exceptions in both subgroups. The isolates sequenced in this study had more homology to some of the recent Pakistan isolates (2005-2008) compared to the former Iranian isolates and in the phylogenetic tree placed closer to new Pakistan isolates than other Iranian isolates. This may indicate appearance of a new sublineage of AI viruses in Iran and Pakistan (fig 1a). Xu *et al* (2007) demonstrated that antigenic diversity among H9N2 viruses from different location in China correspond with phylogenetic relationships (37). Therefore, the observed differences between Iranian H9N2 isolates in phylogenetic analysis reflect their antigenic differences, which cause important challenges in vaccine efficiency.

Most of the amino acids (96%) of Iranian H9N2 virus HAs were highly conservative, whereas the other amino acids showed diverse levels of variability. Certain amino acid substitutions have become fixed in every subgroup, but many are unique substitutions represented in individual viruses. This high

frequency of substitutions in the HA gene of Iranian isolates is a sign of continued selection pressure in the poultry host. It is reasonable that AI vaccination in Iran could enhance this selection pressure and so the rate of the virus adaptation. This could be an important disadvantage of AI vaccination, considering that the AI vaccination cannot prevent occurrence of disease and its economic costs. It should be noted that differences mentioned in this study relate just to the HA gene.

A comprehensive evaluation of differences between Iranian isolates by complete genome sequencing is necessary to understand the probable reassortment of AI H9N2 virus in Iran. The present study highlights the need to continue with influenza virus surveys, to obtain more information that could lead to the prediction and control of future AI H9N2 virus epizooty in Iran.

Acknowledgement

We gratefully acknowledge the excellent technical support provided by Mrs Noroozi. We also thank Lorestan University and University of Tehran for providing part of the financial support.

References

1. Webster RG, Bean WJ, Gorman OT, Chambers TM, Kawaoka Y. Evolution and ecology of influenza A viruses. *Microbiol Rev.* 1992;56:152-179.
2. Fouchier RA, Munster V, Wallensten A, Bestebroer TM, Herfst S, Smith D, Rimmelzwaan GF, Olsen B, Osterhaus AD. Characterization of a novel influenza A virus hemagglutinin subtype (H16) obtained from black-headed gulls. *J Virol.* 2005;79:2814-2822.
3. Nili H, Asasi K. Natural cases and an experimental study of H9N2 avian influenza in commercial broiler chickens of Iran. *Avian Pathol.* 2002; 31: 247-252.
4. Guo YJ, Krauss S, Senne DA, Mo IP, Lo KS, Xiong XP, Norwood M, Shortridge KF, Webster RG, Guan Y. Characterization of the pathogenicity of members of the newly established H9N2 influenza virus lineages in Asia. *Virology.* 2000;267:279-288.

5. Cameron KR, Gregory V, Banks J, Brown IH, Alexander DJ, Hay AJ, Lin YP. H9N2 subtype influenza A viruses in poultry in Pakistan are closely related to the H9N2 viruses responsible for human infection in Hong Kong. *Virology*. 2000;278:36–41.
6. Choi YK, Ozaki H, Webby RJ, Webster RG, Peiris JS, Poon L, Butt C, Leung YH, Guan Y. Continuing evolution of H9N2 influenza viruses in southeastern China. *J. Virol*. 2004; 78:8609–8614.
7. Guan Y, Shortridge KF, Krauss S, Chin PS, Dyrting KC, Ellis TM, Webster RG, Peiris M. H9N2 influenza viruses possessing H5N1-like internal genomes continue to circulate in Southeastern China. *Journal of Virology*. 2000;74:9372–9380.
8. Nili H, Asasi K. Avian influenza (H9N2) outbreak in Iran. *Avian Diseases*. 2003; 47: 828–831.
9. Naeem K, Ullah A, Manvell RJ, Alexander DJ. Avian influenza A subtype H9N2 in poultry in Pakistan. *Veterinary Record*. 1999;145:560.
10. Zhang P, Tang Y, Liu X, Liu W, Zhang X, Liu H, Peng D, Gao S, Wu Y, Zhang L, Lu S. A novel genotype H9N2 influenza virus possessing human H5N1 internal genomes has been circulating in poultry in eastern China since 1998. *J Virol*. 2009;83:8428–8438.
11. Okamatsu M, Sakoda Y, Kishida N, Isoda N, Kida H. Antigenic structure of the hemagglutinin of H9N2 influenza viruses. *Archives of Virology*. 2008;153:2189–2195.
12. Office International des Epizooties. Avian influenza. In: *Manual of Diagnostic Tests and Vaccines for Terrestrial Animals*, fifth ed., Office International des Epizooties, Paris, France, 2008;Chapter 2.7.12.
13. Ghadi S, Bozorgmehri-Fard MH, Karimi V, Tavasoti-Kheiri M, Rafeiei MH, Genetic Analysis of Hemagglutinin Protein of H9N2 Isolated from Live Bird Markets in Tehran Province in 2007–2008. *Iranian J Virol*. 2009;3(3):7–15.
14. Matrosovich M, Zhou N, Kawaoka Y, Webster R. The surface glycoproteins of H5 influenza viruses isolated from humans, chickens, and wild aquatic birds have distinguishable properties. *J Virol*. 1999;73:1146–1155.
15. Kawaoka Y, Webster RG. Sequence requirements for cleavage activation of influenza virus hemagglutinin expressed in mammalian cells. *Proc Natl Acad Sci USA*. 1988;85:324–8.
16. Banks J, Speidel EC, Harris PA, Alexander DJ. Phylogenetic analysis of influenza A viruses of H9 haemagglutinin subtype. *Avian Pathol*. 2000;29:353–60.
17. Liu H, Liu X, Cheng J, Peng D, Jia L, Huang Y. Phylogenetic analysis of the hemagglutinin genes of twenty-six avian influenza viruses of subtype H9N2 isolated from chickens in China during 1996–2001. *Avian Dis*. 2003;47:116–127.
18. Iqbal M, Yaqub T, Reddy K, McCauley JW. Novel genotypes of H9N2 Influenza A viruses isolated from poultry in Pakistan containing NS genes similar to highly pathogenic H7N3 and H5N1 viruses. *PloS ONE*. 2009;4(6):5788.
19. Perk S, Banet-Noach C, Shihmanter E, Pokamnski S, Pirak M, Lipkind M, Panshin A. Genetic characterization of the H9N2 influenza viruses circulated in the poultry population in Israel. *Comp Immunol Microbiol Infect Dis*. 2006;29:207–223.
20. Shahsavandi S, Salmanian AH, Ghorashi SA, Masoudi S, Ebrahimi MM. Evolutionary characterization of hemagglutinin gene of H9N2 influenza viruses isolated from Asia. *Res Vet Sci*. 2012;93(1):234–9.
21. Gambaryan A, Webster R, Matrosovich M. Differences between influenza virus receptors on target cells of duck and chicken. *Arch Virol*. 2002; 147: 1197–1208.
22. Suzuki Y. Sialobiology of influenza: molecular mechanism of host range variation of influenza viruses. *Biol Pharm Bull*. 2005; 28:399–408.
23. Matrosovich M, Gambaryan A, Klenk H. Receptor Specificity of Influenza Viruses and Its Alteration during Interspecies Transmission. In: Klenk H, Matrosovich MN, Stech J, eds. *Avian Influenza: Monographs in virology*, Karger, 2008;27:134–155.
24. Gamblin SJ, Haire LF, Russell RJ, Stevens DJ, Xiao B. The structure and receptor binding properties of the 1918 influenza hemagglutinin. *Science*. 2004;303:1838–1842.
25. Wan H, Perez DR. Amino acid 226 in the hemagglutinin of H9N2 influenza viruses determines cell tropism and replication in human airway epithelial cells. *J Virol*. 2007;81:5181–5191.
26. Aamir UB, Wernery U, Ilyushina N, Webster RG. Characterization of avian H9N2 influenza viruses from United Arab Emirates 2000 to 2003. *Virology*. 2007;361:45–55.
27. Matrosovich MN, Krauss S, Webster RG. H9N2 influenza A viruses from poultry in Asia have human virus-like receptor specificity. *Virology*. 2001;281:156–162.
28. Golender N, Panshin A, Banet-Noach C, Nagar, Pokamnski S, Pirak M, Tendler Y, Davidson I, Garcia M, Perk S. Genetic

characterization of avian influenza viruses isolated in Israel during 2000-2006. *Virus Genes*. 2008;37:289-297.

29. Baigent SJ, McCauley JW. Influenza type A in humans, mammals and birds: determinants of virus virulence, host-range and interspecies transmission. *Bioessays*. 2003;25:657-671.

30. Zhang, P, Tang T, Liu X, Peng D, Liu W, Liu W, Lu S, Liu X. Characterization of H9N2 influenza viruses isolated from vaccinated flocks in an integrated broiler chicken operation in eastern China during a 5 year period (1998-2002). *Journal of General Virology*. 2008;89:3102-3112.

31. Ji K, Jiang WM, Liu L, Chen JM, Chen J, Hou JY, Li JP, Huang BX. Characterization of the hemagglutinin gene of subtype H9 avian influenza viruses isolated in 2007-2009 in China Kang. *J Virol Meth*. 2010;163:186-189.

32. Lee CW, Senne DA, Suarez DL. Effect of vaccine use in the evolution of Mexican lineage H5N2 avian influenza virus. *J Virol*. 2004;78:8372-8381.

33. Rambaut A, Pybus OG, Nelson MI, Viboud C, Taubenberger JK, Holmes EC. The genomic and

epidemiological dynamics of human influenza A virus. *Nature*. 2008;453:615-619.

34. Sun Y, Pu J, Jiang Z, Guan T, Xia Y, Xua Q, Liu L, Ma B, Tian F, Brown EG, Liu J. Genotype evolution and antigenic drift of H9N2 influenza viruses in China from 1994 to 2008. *Vet Microbiol*. 2010; 146 (3-4): 215-225.

35. Karimi V, Bozorgmehri Fard MH, Shahbazzadeh D, Esmaelizad M, Pourbakhsh SA. Sequence analysis and phylogenetic study of hemagglutinin gene of H9N2 subtype of avian influenza virus isolated during 1998-2002 in Iran, *Iranian Biomedical J*. 2004;8(4):167-172.

36. Homayounimehr AR, Dadras H, Shoushtari A, Pourbakhsh SA. Sequence and phylogenetic analysis of the hemagglutinin genes of H9N2 avian influenza viruses isolated from commercial chickens in Iran. *Trop Anim Health Prod*. 2010;42:1291-1297.

37. Xu KM, Li KS, Smith GJD, Li JW, Tai H, Zhang JX, Webster RG, Peiris JSM, Chen H, Guan Y. Evolution and molecular epidemiology of H9N2 influenza A viruses from quail in southern China: 2000 to 2005. *J Virol*. 2007;81:2635-2645.